



## Role of Lactobacillus Probiotics in Gut Health and Food Safety: Mechanisms for Contaminant Reduction and Applications in Food Products

Mohsin Gulzar Barq<sup>a\*</sup>, Amna Razzaq<sup>a</sup>, Manal Yasin<sup>a</sup>, Khadija Nawal<sup>a</sup>, Tayyaba Qureshi<sup>a</sup>

a. University of Central Punjab, Avenue 1 Khayaban-e-Jinnah, Pir Mansur Johar Town, Lahore, Punjab

**Abstract:** Lactobacillus probiotics, live bacteria that confer health benefits when administered in adequate amounts, have gained widespread acceptance, particularly due to their potential to influence and enhance general health by modulating the gut microbiota. In recent years, probiotics have been highlighted for their potential therapeutic applications in preventing and mitigating various diseases. Among these, Lactobacillus spp. represent one of the most extensively studied and utilized probiotic strains. This review explores the role of Lactobacillus in maintaining gut microbiota balance, producing bioactive metabolites such as short-chain fatty acids (SCFAs) and bacteriocins, and its involvement in food safety through the reduction of contaminants. Specifically, Lactobacillus spp. contribute to food safety by adsorbing, enzymatically degrading, or neutralizing harmful substances, including aflatoxins and heavy metals, thereby reducing their bioavailability and toxicity. The paper also discusses the practical applications of Lactobacillus-based probiotics in the food industry and the ongoing efforts to address associated safety concerns. Lactic acid bacteria (LAB) and probiotics provide a natural, non-chemical approach to food preservation by inhibiting pathogenic microorganisms and extending shelf life. Their incorporation into food systems promotes a safer and more sustainable food supply. However, further research is required to optimize their applications and ensure their efficacy in diverse food matrices.

**Key words:** Lactic acid bacteria, Probiotics, Lactobacillus casei, Food contaminants, Streptococcus thermophiles, Food industry

### 1. Introduction

Lactic acid bacteria (LAB) are a group of beneficial microorganisms widely known for their ability to ferment sugars into lactic acid. LAB belongs predominantly to the phylum Firmicutes and is classified as Gram-positive, non-sporulating, and either rod or cocci-shaped ([Miranda et al., 2021](#)). They thrive in anaerobic or microaerophilic conditions, meaning they grow best in environments with little or no oxygen ([Bintsis, 2018](#)). LAB is important in food preservation and contributes to fermented products' flavor, texture, and safety. By producing lactic acid, they lower the pH of their environment, inhibiting the growth of spoilage microorganisms and pathogens. This acidification process is crucial for the production of yogurt, cheese, sauerkraut, kimchi, pickles, sourdough bread, and various fermented beverages. LAB is also significant in producing probiotics, contributing to gut health by promoting a balanced intestinal microbiota ([Pérez-Rivero & López-Gómez, 2023](#)). Lactobacillus strains are amongst the most studied probiotics, given their diverse species and strains with distinct characteristics and functions. The beneficial effects of these strains include immune function reinforcement and decreased incidence of gastrointestinal illnesses, in addition to the beneficial impacts on digestion and nutrient absorption ([Soomro, 2002](#)). One of the key ways that Lactobacillus demonstrates its probiotic effects is via its promotion of healthy microbiota in the gut ([Soomro, 2002](#)). Lactobacillus uses probiotic benefits to influence and modulate the gut microbiota, leading to beneficial impacts on human health, such as inhibition of colonization and growth of pathogens in the gastrointestinal tract, production of antimicrobial compounds that have proven to be effective, promotion of cytokine production, and improvement of intestinal barrier integrity. The gut microbiota composition and function may be disrupted in connection with various gastrointestinal conditions and systemic diseases ([Collado et al., 2007](#)). Probiotics, specifically Lactobacillus probiotics, have been studied intensively for their effect on influencing the gut microbiota and resulting health benefits to the host. This review focuses on the diverse habitats in which

[Received] 21 Oct 2024; Accepted 11 Dec 2024; Published 22 Dec 2024]

Finesse Publishing stays neutral about jurisdictional claims published maps




Attribution 4.0 International (CC BY 4.0)

Corresponding email: [mohsin.barq@yahoo.com](mailto:mohsin.barq@yahoo.com) (Mohsin Gulzar Barq)

DOI: 10.61363/mapd4e52

*Lactobacillus* species can be found, including the human gastrointestinal tract, dairy products, and fermented products. Probiotics refer to live bacteria that have health-promoting properties to the host, and *Lactobacillus* is a significant probiotic genus. The beneficial potential of probiotics for health maintenance has been largely studied. *Lactobacillus* species exhibit several important characteristics that make them attractive probiotic candidates. Firstly, *Lactobacillus* species are present in fermented foods and as dietary supplements and are considered safe for human consumption.

They are a normal part of the human gastrointestinal microbiota, and their presence is associated with a healthy digestive system (Kongo, 2013). Secondly, the survival and functionality of *Lactobacillus* species are ensured by their ability to withstand the extremely acidic environment of the stomach and reach the intestines in a viable form. The resistance of these bacteria is attributed to their strong cell walls and ability to tolerate acid. Thirdly, *Lactobacillus* species have the remarkable ability to attach to intestinal epithelial cells, which allows them to colonize and improve the gut environment (Khatoon et al., 2023). This allows *Lactobacillus* species to outcompete pathogenic bacteria and prevent their colonization. Food contamination remains a critical issue globally, arising from microbial agents such as *Escherichia coli*, *Salmonella*, and *Listeria*, as well as chemical contaminants like heavy metals, pesticides, and industrial toxins (Hussain & Gooneratne, 2017). These contaminants compromise food safety and have severe public health implications. The consumption of contaminated food is one of the leading causes of foodborne diseases, which affect millions globally each year, leading to significant morbidity, mortality, and economic losses (DiStefano, 2014). Food contaminants can enter the food supply from farm to fork, through sources like air, water, soil, packaging materials, processing equipment, and intentional adulteration (Hussain & Gooneratne, 2017). With the global impact of foodborne diseases rising, preventative strategies are increasingly focused on reducing contamination risks (Rather et al., 2017).



**Figure 1.** This shows the general taxonomy and key characteristics of Lactic acid bacteria

Traditional food preservation methods, though effective, have limitations in fully addressing these challenges. As a result, LAB and probiotics have emerged as promising tools in combating microbial contamination and degrading certain harmful chemicals in food. This review aims to explore the reduction potential of LAB and probiotics in eliminating food contaminants and to highlight their applications across various food products. By understanding their mechanisms of action and the diverse ways they can be integrated into the food supply chain, this review will underscore the role of these microorganisms in improving food safety and contributing



to public health. Finally, the wide variety of metabolic activities of *Lactobacillus* species, including the enzymes and anti-microbial compounds that facilitate their antibacterial and immune-modulatory activities ([Mezaini & Bouras, 2013](#)). These features collectively give *Lactobacillus* species high therapeutic potential as probiotics, and, as such, are highly regarded for their capacity to promote gut health and well-being.

## 2. Contaminants in food

Food contaminants present a serious global public health challenge, with biological, chemical, and physical contaminants contributing to various health risks. Preventing contamination at all stages of the food supply chain, farming, processing, packaging, and storage, is essential to ensure food safety. Regulatory bodies, such as the FDA (Food and Drug Administration) and WHO (World Health Organization), play a crucial role in monitoring and enforcing food safety standards to mitigate these risks. Moreover, consumers must practice safe food handling, cooking, and storage techniques to reduce the risk of contaminant exposure further. (Table 1)

**Table 1.** This shows the contamination rate in different food products

Products	Contamination Rate	Reference
Ricotta	86.2 %	<a href="#">(Béjaoui et al., 2022)</a>
Cheese	69.6 %	<a href="#">(Béjaoui et al., 2022; Serpe et al., 1999)</a>
Raw milk	72.5 %	<a href="#">(Béjaoui et al., 2022)</a>

### 2.1. Contamination by bacteria

Threats due to bacterial species are called Brucellosis, which causes a foodborne disease in humans, and it may be mild or chronic depending on the severity of the infection. It mainly spreads through animals or by consuming infected meat or products. Dairy products, mainly ricotta, cheese, or raw milk, may be contaminated by *Brucella* spp. like *B. suis*, *B. melitensis*, and *B. abortus*. The well-affected areas due to Brucellosis are rural areas due to touching infected animals directly, while people living in urban areas may ingest unpasteurized dairy products. The ingestion of contaminated products may be fatal and cause meningitis, endocarditis, or hepatitis diseases ([Béjaoui et al., 2022; Serpe et al., 1999](#)). Illness caused by *Escherichia* produces Shiga toxin, which is present in fermented dairy products. Thus, causing foodborne diseases, especially in products like Kashk and Doogh ([Dehkordi et al., 2014](#)). From bulk farm milk, market milk, fresh soft cheese, kareish cheese, and ice cream, *Salmonella* spp. Can contaminate products, mostly *S. typhimurium*. *S. enteritidis* and *S. infantis* present ([Omar et al., 2018](#)). Two common species, i.e., *S. Typhimurium* & *S. enteritidis*, pathogens of milk and its products, mostly in raw sheep milk ([Rostami et al., 2014](#)).


According to several studies, in dairy products like cheese and milk (liquid), gram-negative bacteria, i.e., coliforms, are mainly present as a contaminant ([Hervert et al., 2017](#)). *Paenibacillus* or *Bacillus* causes spoilage in milk, and it is tough to remove because it possesses characteristics of endospore-forming bacteria ([Huck et al., 2007](#)). Other pathogens that affect milk are *Brucella abortus*, *Campylobacter*, *Clostridium*, *E. coli*, *Salmonella*, *Mycobacteria*, and *Streptococcus agalactiae* ([Msalya, 2017](#)). *Listeria monocytogenes* also contaminates milk products. Cold raw milk favors the growth of these bacteria ([Kabuki et al., 2004](#)). Besides the presence of gram-negative bacteria, gram-positive aerobic bacteria are also present, like *Staphylococcus intermedius*, *Staphylococcus aureus*, *Streptococcus agalactiae*, and *Staphylococcus epidermis* ([Mdegela et al., 2004](#)). Yogurt also becomes contaminated by these gram-negative bacteria, i.e., Enterobacteriaceae, or also through *E. coli*. Through EB testing, we can know whether yogurt is hygienic or not ([Hervert et al., 2017](#)). One of the common spoilage microorganisms, *Enterobacter cloacae*, was likely to survive in cow milk. In soymilk, *Pseudomonas paucimobilis* is present and survives in yogurt ([Canganella et al., 1999](#)). Anaerobic facultative spoilage microbe, i.e., *Listeria monocytogenes*, is present and causes the disease Listeriosis in humans, which greatly affects the liver & spleen ([Yang & Yoon, 2022](#)). Gram-positive bacteria, i.e., *Listeria monocytogenes*, contaminate the product, especially in fresh cheese processing plants, and harm human health if consumed. Also, *Pseudomonas*, *Salmonella* spp, *E. coli*, and *Bacillus cereus* are present ([Al-Gamal et al., 2019](#)). Cheese production from raw milk was mostly used, and common spoilage microbes are Verotoxigenic *E. coli*, *S. aureus*, and *Salmonella* (Table 2) ([Costanzo et al., 2020](#)).

**Table 2.** Food-borne bacteria and lactic acid bacteria against them

Pathogenic Bacteria	How LAB fights against them
<i>Listeria monocytogenes</i>	LAB produces bacteriocins like nisin and organic acids that lower the pH, making it difficult for <i>Listeria</i> to survive.
<i>Salmonella spp.</i>	They create organic acids and compete for nutrients, effectively lowering the pH and limiting <i>Salmonella</i> 's growth.
<i>Escherichia coli</i> O157	LAB produces organic acids, hydrogen peroxide, and bacteriocins, all of which help to combat <i>E. coli</i> .
<i>Staphylococcus aureus</i>	They generate reuterin, along with organic acids and bacteriocins, to inhibit <i>Staphylococcus</i> growth.
<i>Clostridium perfringens</i>	LAB produces organic acids and bacteriocins that help reduce the presence of <i>Clostridium</i> in food.
<i>Bacillus cereus</i>	By producing bacteriocins and lowering the pH, LAB can inhibit the growth and spore formation of <i>Bacillus</i> .

**2.2. Contamination by Fungi**

In fermented products, highly toxigenic molds are present, which represent mycotoxins. Airborne infection, improper use of equipment, or handling refers to a source of contamination in products (Pei et al., 2021). It can be affected by Aflatoxin, especially when milk is in liquid form, which is produced by the species *Aspergillus flavus* and causes a contagious effect on the health of humans. Dairy products may be affected by it even after passing through pasteurization. Collection of milk safely, storage, and temperature level are important in protecting products from MI(AFM1). (Figure 2)



**Figure 2.** Types of aflatoxin and their impact on the Human Body

However, due to its growth rate at low temperatures, the fermentation process between product ingredients like lactose and sucrose, hydrolysis of protein, or formation of lactic acid in products, these characteristics may lead to the formation of aflatoxin. Another species, *Aspergillus parasiticus*, also has a carcinogenic effect on the liver, kidney, and its functioning. Different applications are applied to remove this, but they cannot give 100 % accuracy. Aflatoxin compound M1(AFM1) causes dangerous effects on the health of humans (Bayat et al., 2021).



Aflatoxin sometimes produces mycotoxin, which is referred to as ochratoxin, which has hazardous effects not only on humans but also on animals ([Onoharigho et al., 2022](#)). Yeast and mucor are also present in it ([Bastam et al., 2021](#)). Some other species are also important according to a study conducted in Isfahan, which shows that Acremonium & Cladosporium are present, which contaminate milk and its related products (Table 3).

**Table 3.** Isolates of fungi cause contamination in different prepared food products.

Product	Fungal Isolates	Contamination Rate	Reference
Peanut butter	<i>Helmintosporium</i>	47.44%	<a href="#">(Zamblé et al., 2016)</a>
	<i>Penicillium</i>	41.54%	
	<i>Aspergillus sp</i>	20.22-51.65%	
Cooking butter	<i>Aspergillus</i>	2.8%	<a href="#">(Ismail &amp; Sabreen, 2001)</a>
	<i>Geotrichum</i>	78%	
	<i>Penicillium</i>	3.9%	
	<i>Mycosphaerella</i>	1.4%	

It may be affected by spoilage due to yeast, which serves as the topmost contaminant in it. As yogurt is stored at low temperatures for its proper formation and its acidic environment, yeast is more likely to grow, especially *Saccharomyces* or *Candida kluyveromyces* ([Al-Gamal et al., 2019](#)). Common spoilage microorganisms that are involved in the spoilage of yogurt are *Penicillium monilia*, *Alternaria*, *Rhizopus*, *Micelia sterilis*, and *Cladosporium*. Contamination through the air is very common, and it makes products spoil, especially through *Penicillium* or *Clostridium*. For this purpose, certain food additives like Potassium sorbate are added to it. Bio preservatives enhance product quality as well as increase its shelf life. Different LABs are used to make fruits, vegetables, or dairy products free from contaminants ([Buehler et al., 2018](#)). Contaminants that affect butter, mainly molds or yeast, change the color of the product and also make its flavor bitter ([Ei-Shafe et al., 2017](#)). Butter is mainly spoiled by *Penicillium spp.* and *Cladosporiodes spp.* Most importantly affected by *Geotrichum candidum*. About 78% of it spoils cooking butter ([Ismail & Sabreen, 2001](#)). Yeast caused spoilage in cheese. Moreover, it changes its texture, smell, color, and taste, etc ([Ei-Shafe et al., 2017](#)). Depending upon the type of cheese, various microorganisms spoil it, such as soft cheese. *Aspergillus*, *Mucor*, and *Penicillium* cause spoilage in this product. Cheddar cheese was commonly affected by *Aspergillus*, *Penicillium*, or *Fusarium* ( Table 4) ([Ismail & Sabreen, 2001](#)).

**Table 4.** Shows the different types of fungal species that spoil the different types of cheeses.


Product Name	Commonly Found In The Country	Common Spoilage Microorganism	Reference
Processed Cheese	Egypt	<i>Aspergillus</i>	<a href="#">(Ismail &amp; Sabreen, 2001)</a>
		<i>Penicillium</i>	
Hard Cheese	Egypt	<i>Rhizopus</i>	<a href="#">(Ismail &amp; Sabreen, 2001)</a>
		<i>Penicillium</i>	
		<i>Aspergillus</i>	
		<i>Geotrichum</i>	
		<i>Candida</i>	
		Other yeast	
Cheese	New Zealand & Australia	<i>Penicillium commune</i>	<a href="#">(Ismail &amp; Sabreen, 2001)</a>
		<i>P. roquefortii</i>	
		<i>P. chrysogenum</i>	
		<i>P. expansum</i>	
		<i>P. solitum</i>	
		<i>P. viridicatum</i>	
Packaged Cheese	Europe	<i>P. brevicompactum</i>	<a href="#">(Ismail &amp; Sabreen, 2001)</a>
		<i>P. commune</i>	
		<i>P. verrucosum</i>	
		<i>P. solitum</i>	
		<i>P. roquefortii</i>	

		<i>P. nalgiovensis</i>	
<b>Cheddar Cheese</b>	Egypt	<i>Aspergillus</i> <i>Penicillium</i>	(Ismail & Sabreen, 2001)
<b>Soft Cheese</b>	Egypt	<i>Candida</i> <i>Penicillium</i>	(Ismail & Sabreen, 2001)

**3. Reduction mechanism of bacterial contaminants in food**

The general aim of shelf-life extension is to keep foods safe and stable. Generally, this can be achieved by controlling the growth of spoilage microorganisms and pathogenic bacteria. In treating and controlling pathogenic growth, an antimicrobial agent, such as nisin, or two or more antimicrobial agents could be used synergistically against the target organism. The majority of reported foodborne illnesses are caused by pathogenic bacteria such as *Campylobacter jejuni*, *S. aureus*, *L. monocytogenes*, *E. coli*, and *Salmonella* spp. Consumer demand for minimally processed foods has obligated the food industry to search for new methods of ensuring food safety. It can no longer rely on traditional heat treatment methods to create microbiologically safe foods. On the other hand, fruits and vegetables have been found to harbor pathogenic bacteria. LAB may enhance the nutritional quality of food and also contribute to intestinal health due to the production of antimicrobial agents. There are various mechanisms for foodborne pathogens prevention and elimination of food spoilage bacteria, like the production of antimicrobial substances that may prevent adherence of pathogens to epithelial and mucosal surfaces. LAB produces different compounds that help to reduce the contamination rate by bacterial species in food. It includes organic acids, bacteriocin production, hydrogen peroxide production, competitive exclusion, and Stimulation of Host Defenses (in fermented foods).

LABs are also able to inhibit the fungi (molds) responsible for food contamination and mycotoxin production. LABs contain bacteriocin-like substances and produce organic acids with typical fungistatic and fungicidal properties, and inhibit fungi and yeast such as *Aspergillus versicolor*, *Penicillium expansum*, *Fusarium culmorum*, *Candida parapsilosis*, *Aspergillus niger*, and *Penicillium chrysogenum*. Some species of several genera of fungi produce mycotoxins, which are *Aspergillus*, *Fusarium*, *Penicillium*, and *Alternaria*. These fungi genera produce different Mycotoxins, including aflatoxins, fumonisins, ochratoxin, patulin, tricothecenes, and zearalenone, which are derived as secondary metabolites. These metabolites have potential carcinogenic, teratogenic, immunotoxic, neurotoxic, hepatotoxic, and nephrotoxic effects. Among approximately 400 compounds identified as mycotoxins, 30 of them are considered dangerous to human health (Bayat et al., 2021). Thus, myco-toxin contamination should be controlled either by preventing their formation or by detoxification. The control of mycotoxins is more difficult because the normal cooking method cannot destroy all of them. Therefore, food processing techniques are necessary to remove mycotoxins. LABs (*Lactocaseibacillus casei* and *Limosilactobacillus reuteri*) are known to efficiently bind toxins. Such toxins include AFs (aflatoxin) in aqueous solutions. Thus, the control of these microbes using LAB is a natural way of food preservation. (Figure 3).



**Figure 3.** Mechanism of reduction of contaminants with *Lactobacillus* and their role in food products.



### 3.1. Production of organic acid

Lactic acid bacteria (LAB) play a crucial role in keeping food safe using several natural mechanisms. They produce compounds like lactic acid, acetic acid, and reuterin, which lower the pH of the food, making it harder for harmful bacteria to grow. LAB also competes with pathogens for nutrients, quickly consuming what is available and creating an acidic environment that further stops the growth of harmful microbes. Additionally, they can break down glycerol, releasing antimicrobial substances that help preserve food. Beyond just keeping food safe, LAB also improves the taste and extends the shelf life, acting as a natural defense against spoilage. The wide variety of LAB strains brings an even broader range of protective benefits, making them highly effective in preventing foodborne illness ([Gänzle et al., 2024](#)).

Lactic acid bacteria (LAB) play an important role in protecting food from various harmful food-borne bacteria such as *Listeria monocytogenes* and *Staphylococcus aureus*. LAB also excels at competing for nutrients, quickly using up available resources and producing organic acids that inhibit the growth of harmful bacteria like *Salmonella* spp. and *Escherichia coli* O157:H7 ([Cálix-Lara et al., 2014](#)). Additionally, they release hydrogen peroxide and other antimicrobial compounds that further strengthen their protective effects. The lowered pH disrupts bacterial cell membranes, inhibits enzyme activity, and reduces the growth of spoilage organisms and pathogens like *Escherichia coli*, *Salmonella*, and *Listeria*.

### 3.2. Production of bacteriocin

These LAB-produced bacteriocins are antimicrobial peptides that can inhibit many other bacteria, such as pathogens and spoilage organisms. These include *Lactobacillus*, *Lactococcus*, *Streptococcus*, and *Enterococcus*, among the various species of LAB, which spontaneously produce bacteriocins as part of a defense mechanism for competition in the gut or food products. Of specific interest are these bacteriocins due to their potential application as natural preservatives for foods and as alternatives to antibiotics ([Parada et al., 2007](#)). There are two major classes of bacteriocin. Class I bacteriocin is called Lactococcin are small heat-stable peptide that doesn't require post-translation modifications. When LAB produces this type of bacteriocin, it directly targets inhibiting the peptidoglycan synthesis of other contaminants present in food. In this way, contamination reduction can be easily performed. The other class of bacteriocin is Class II bacteriocin, which is called Lantibiotics, which function by directly destabilizing the cytoplasmic membrane of the contaminants present in food with the help of the creation of pores that allow bacteriocin molecules to enter into the bacteria and the cells death. In this way, microbial contamination can easily be reduced with the help of different bacteriocins (Table 5) ([Ibrahim et al., 2021](#)).

**Table 5.** Mechanisms of lactic acid bacteria in enhancing food safety and quality

Mechanism	Description
Production of Antimicrobial Compounds	LAB naturally produces substances like lactic acid, acetic acid, and reuterin, which lower the pH of food and help inhibit the growth of harmful bacteria.
Competitive Exclusion	LABs compete with harmful bacteria by quickly using up available nutrients, producing acids that create an environment where pathogens struggle to survive.
Utilization of Glycerol	LAB can break down glycerol, releasing antimicrobial compounds in the process. This helps to preserve food and keep it free of harmful microbes.
Impact on Food Quality and Safety	The metabolic activities of LAB not only help prevent the growth of pathogens but also enhance the taste and shelf life of food products.
Diversity of LAB	The metabolic activities of LAB not only help prevent the growth of pathogens but also enhance the taste and shelf life of food products.

#### **4. Enhancing Food Products with Lactic Acid Bacteria**

Lactic acid bacteria (LAB), especially *Lactobacillus*, play a big role in making food healthier, tastier, and safer. LAB is essential for the process of fermentation, where they turn sugars into lactic acid. This gives food a unique flavor and texture and naturally preserves it by lowering the pH, which stops harmful bacteria from growing. This process helps extend the shelf life of many foods while keeping them safe to eat. LAB helps your body to absorb nutrients more easily. For instance, dairy products break down lactose, which means people who are lactose intolerant can enjoy milk and yogurt without discomfort. So, they make these foods more accessible and easier to digest ([Al-Gamal et al., 2019](#)).

*Lactobacillus* strains are known for their probiotic benefits. They help keep the balance of good bacteria in your gut, which is important for digestion and immune health. Adding these probiotics to foods like yogurt or fermented veggies offers extra health perks. Additionally, LAB produces natural substances that protect food from spoilage. By creating antimicrobial compounds, they prevent the growth of harmful bacteria and fungi, making food safer and helping to avoid foodborne illnesses. These helpful bacteria also make food taste and feel better. The flavors in fermented foods like yogurt, cheese, and pickles give them their signature tang and texture that people love. These probiotics support digestion, strengthen immunity, and even have anti-inflammatory effects, making them a favorite for health-conscious consumers ([Zhang et al., 2018](#)).

#### **5. The Role of Lactobacillus Strains in Health and Disease Prevention**

*Lactobacillus* strains, that is a key group of probiotics, have demonstrated significant health benefits, particularly in promoting gut health and preventing various diseases.

##### **5.1. Lactobacillus salivarius CECT 5713 and Digestive Health**

This strain plays a crucial role in preventing digestive issues, such as indigestion. Studies have shown that infants who were fed formula containing *Lactobacillus salivarius* for up to six months experienced a notable increase in fecal lactobacilli ([Maldonado et al., 2010](#)). This boost in beneficial gut bacteria helps maintain a balanced microbiome, which is essential for proper digestion and overall gut health.

##### **5.2. Lactobacillus rhamnosus GG and Infection Prevention**

*Lactobacillus rhamnosus* GG is widely used in fermented milk products and has been shown to reduce the risk of respiratory and gastrointestinal infections prevalent in developing countries, affecting 5-44% of the population. Regular consumption of products containing this probiotic strain can significantly lower the incidence of such infections by supporting the immune system and maintaining gut integrity ([Hojsak et al., 2010](#)).

##### **5.3. Lactobacillus acidophilus and Lactose Intolerance**

*Lactobacillus acidophilus* is particularly effective in enhancing lactose utilization in individuals who are lactose intolerant. These individuals lack lactase, the enzyme responsible for breaking down lactose, often leading to symptoms like diarrhea and abdominal discomfort. By consuming *L. acidophilus*-rich products, lactose digestion is improved ([Kim & Gilliland, 1983](#)).

##### **5.4. Yogurt and Gut Health**

*Lactobacillus casei* is commonly used to treat diarrhea, especially in children. Fermented yogurt containing *L. casei* has been found to significantly improve gut health by restoring the balance of beneficial bacteria in the gut, thus reducing the severity and duration of diarrhea ([Pereg et al., 2005](#)).

##### **5.5. Gut Microbiota and Protection Against Harmful Bacteria**

Probiotics such as *Lactobacillus casei*, *L. acidophilus*, and *Bifidobacterium* strains, naturally present in the human intestine, contribute to maintaining a healthy gut environment. These strains produce organic acids and bacteriocins, which inhibit the growth of harmful bacteria by creating an inhospitable environment for pathogens. This antimicrobial activity is essential for promoting a healthy gut and preventing infections (Figure 4) ([Mazahreh & Ershidat, 2009](#)).


##### **5.6. Fermented Beverages: Their Role in Health and Production**

Fermented beverages have gained popularity due to their health benefits and are commonly produced by introducing lactic acid bacteria (LAB) or yeast during fermentation. These beverages not only improve sensory



qualities like flavor and texture but also serve as a source of probiotics that enhance gut health. Lactic acid beverages are primarily prepared using milk or dairy products. Key bacterial strains involved in their production include *Lactobacillus acidophilus*, *Lactobacillus rhamnosus*, and *Bifidobacterium animalis*. Research indicates that these beverages contain around 51% lactic acid, contributing to their distinct taste and health benefits ([Almeida et al., 2009](#)). Factors such as the physical, sensory, and chemical properties of the beverage are crucial in optimizing production and enhancing quality. These drinks are a rich source of live microorganisms that can colonize the human gut, helping prevent diseases by promoting a balanced microbiome and improving digestive health. Dairy-based fermented beverages typically consist of a mixture of whey and milk (usually from buffalo or cows).

Strains like *Lactobacillus bulgaricus* and *Lactobacillus acidophilus* are commonly used in their production. Whey, a by-product of cheese production, has a reputation for promoting health due to its high nutrient content, including calcium (Ca), iron (Fe), magnesium (Mg), and phosphorus (P), while maintaining low cholesterol levels. However, improper disposal of whey from the dairy industry can pose environmental hazards. These dairy beverages are particularly beneficial for treating infections caused by gut microbiome imbalances, further highlighting their probiotic properties ([da Silva et al., 2020](#)). Recent studies have explored the combination of fruit juices and whey to create probiotic beverages ([Worku et al., 2019](#)). For example, pineapple juice mixed with whey in a 65:35 ratio, inoculated with 1% *Lactobacillus acidophilus*, has been shown to promote the growth of beneficial gut bacteria. Additionally, *Lactobacillus plantarum* is known to enhance gut health when used in fermented fruit beverages. A notable example is the Cornelian cherry, which possesses unique antimicrobial and antimalarial properties, making it a valuable ingredient in probiotic drinks ([Mantzourani et al., 2018](#)).



**Figure 4.** Role of Lactobacillus in milk products and enhancing human health.

## 6. Industrial applications of LAB

LABs have a large number of modern applications because of their exceptional metabolic properties, security, and capacity to flourish in different conditions. In the emerging field of biotechnology, LABs have arisen as an overlooked yet truly great individual using their flexible capacities across modern applications. From the development of mature food varieties to the pharmaceutical industry, and then some, to the textile industry, these have proved to be excellent microorganisms in every industrial application. They are broadly utilized as starter societies in the maturation of dairy items like cheddar, yogurt, and aged milk, as well as in meat, fish, organic products, vegetables, and cereal products ([Bintsis, 2018](#)). LAB contributes altogether to the flavor, surface, and healthy benefits of aged food varieties, acting as assistant societies and speeding up cheddar (cheese) development. Additionally, they produce bacteriocins and antifungal mixtures, prompting the use of bio-defensive societies in food preservation ([Miranda et al., 2021](#); [Zapašnik et al., 2022](#)).

## **6.1. Commercial use of LAB in the food industry**

Many researchers have reported the good potential of LAB on humans or animals when they consume fermented food, which then shows good benefits on their health. The fermentation processes, biosynthetic ability, and metabolic activities are the key elements that work with the use of LAB as microbial starters for creating, delivering, and expanding explicit gainful mixtures in aged (fermented) food ([López & Spano, 2023](#)). LAB ferments carbohydrates to Lactic acid, so it is mostly used in the fermented food industry. There are several commercial uses of LAB in specific food industries in different processes.

### **6.1.1. Versatility in food fermentation**

LAB plays a crucial role in dairy and non-dairy food products, contributing to their fermentation, flavor, and potential health benefits. These bacteria, commonly found in fermented dairy foods, include genera like *Lactobacillus*, *Lactococcus*, etc. LAB strains isolated from dairy environments have shown protection from low pH, bile salts, and recreated gastric and gastrointestinal circumstances, along with beneficial enzymatic activities and genes that indicate probiotic potential ([Coelho et al., 2022](#); [Colombo et al., 2018](#)). LAB in dairy products can enhance lactose digestion, stimulate the immune system, and even stop the growth rate of pathogens like *Staphylococcus aureus*, *Escherichia coli*, and *Salmonella typhimurium* ([Ağagündüz et al., 2021](#)). Their presence in dairy products contributes to the rich flavor, texture, and potential health-promoting effects associated with these products ([Coelho et al., 2022](#)). LAB additionally has well-being-advancing properties, making them significant as probiotic societies in the food industry ([Zapašnik et al., 2022](#)). Furthermore, LAB exhibits antimicrobial action against foodborne microorganisms, adding to sanitation by hindering the development of destructive microbes, yeasts, and growths through the creation of natural acids like lactic acid, acetic acid, and propionic acid ([Wang et al., 2021](#)). Their capacity to kill mycotoxins and produce metabolites, for example, bacteriocins, further upgrades their role in food preservation ([Zapašnik et al., 2022](#)). Generally, LAB's metabolic qualities and antimicrobial properties make them fundamental for further developing the taste, surface, security, and shelf life of different food items in the business.

### **6.1.2. Yogurt and Cheese**

One of the most important products of the dairy industry is yogurt, which is the best example of a fermented product in which LAB is used, and gives us the best form of yogurt ([Chen et al., 2017](#)). The rate of consumption of yogurt is very high in the whole world. It helps to improve our gut health. It is delivered through the fermentation of milk by unambiguous types of microorganisms, principally *Lactobacillus bulgaricus* and *Streptococcus thermophiles*, albeit extra probiotic microbes may likewise be incorporated, relying upon the ideal qualities of the yogurt ([Colombo et al., 2018](#); [Mazahreh & Ershidat, 2009](#)). During fermentation, these microbes convert lactose, the regular sugar present in milk, into lactic acid. This fermentation cycle makes the milk proteins coagulate, bringing about the thick, smooth surface attribute of yogurt. The aging likewise adds to the tart flavor and particular smell of yogurt ([Bintsis, 2018](#)). Yogurt is valued for its taste and texture as well as for its various medical advantages. It is a rich wellspring of fundamental supplements like calcium, protein, nutrients B6 and B12, riboflavin, and potassium. National Yogurt Association rules characterize the dynamic culture yogurt as a result that contains live (LAB) microbes in a sum of > 10<sup>8</sup> cells/g at the end of the production ([Mazahreh & Ershidat, 2009](#)).

Moreover, the live microscopic organisms' societies present in yogurt, frequently referred to as probiotics, can assist with keeping a good balance of stomach microorganisms, help in processing, support resistance, and possibly mitigate the side effects of lactose intolerance. Limited research on humans concerning the gut-brain axis focuses on LAB, which is linked to brain functioning, psychology, and immunology. For instance, fermented milk products containing specific LAB strains improved cognitive performance and modulated immune responses in different populations. In milk fermentation, the *Lactobacillus casei* DN-114001 strain of LAB, which is mostly present in yogurt, helps to improve our immune system ([Marcos et al., 2004](#)). While consumption of unpasteurized milk increased *Lactobacillus* levels and influenced microbiome profile, no direct link between LAB amount and mental or psychological measures has been established, indicating a need for further research on the direct effects of dairy LAB on the gut-brain axis (Table 6) ([Ağagündüz et al., 2021](#); [Butler et al., 2020](#)).

**Table 6.** Shows the different parameters on which Lactic acid bacteria isolates grow and the number of bacterial isolates present in different dairy food products.



Industrial product	LAB strains	Salt concentration, temperature, and pH	Bacterial amount in the end product	References
<b>Fermented dairy products</b>				
<b>Yogurt</b>	<i>Lactobacillus bulgaricus</i> <i>Streptococcus thermophiles</i> <i>Lactobacillus Acidophilus</i> <i>Bifidobacterium</i> <i>Lactobacillus casei</i>	Tolerating environments with a low pH 2% or eventually 4% of salt, 42°C is the ideal growth temperature for a starter culture	About > 10 <sup>8</sup> cells/g	( <a href="#">Colombo et al., 2018</a> ; <a href="#">Hosken et al., 2023</a> ; <a href="#">Kongo, 2013</a> ; <a href="#">Mazahreh &amp; Ershidat, 2009</a> )
<b>Cheese</b>	<i>Lactococcus lactis</i> <i>S. salivarius</i> subsp. <i>Thermophilus</i> <i>Lb.helveticus</i> , <i>Lb.delbrueckii</i> subsp <i>Delbrueckii</i> subsp. <i>bulgaricus</i> <i>Ln. lactic</i> <i>Ln. cremoris</i> <i>S. salivarius</i> subsp. <i>Thermophilus</i>	The temperature for cheese ripening is 15°C pH is about neutral, high activity of water Good source of nutritional The mesophilic starter culture temperature is 30 °C and 40 and 45 °C is for Thermo tolerant species	2.3×10 <sup>7</sup> CFU per gram.	( <a href="#">Coelho et al., 2022</a> ; <a href="#">Hosken et al., 2023</a> ; <a href="#">Kongo, 2013</a> ; <a href="#">Nicosia et al., 2023</a> )
<b>Kefir</b>	<b>LAB strains:</b> <i>Lactobacillus kefiranofaciens</i> , <i>Lactobacillus kefir</i> , <i>Lactobacillus plantarum</i> , <i>Lactococcus lactis</i> ssp. <i>lactis</i> , <i>Kluyveromyces marxianus</i> , <i>Lb. acidophilus</i> <i>Lb. parakefir</i> <i>Lb. casei</i> <i>Lb. rhamnosus</i> , <i>Lb. fructivorans</i> , <i>Lb. hilgardii</i> <b>Yeast strains:</b> <i>Zygosaccharomyces</i> sp. <i>Candida lipolytica</i> , <i>C.holmii</i> <i>Saccharomyces cerevisiae</i> <i>S. fragilis</i> <i>S. lactis</i>	The temperature for LAB Strains to ferment kefir is 37°C	10 <sup>8</sup> CFU/mL of lactobacilli and lactococci and 10 <sup>5</sup> cfu/mL of yeasts Acetic acid bacteria (10 <sup>5</sup> CFU/g)	( <a href="#">Arslan, 2015</a> ; <a href="#">Prado et al., 2015</a> ; <a href="#">Tan et al., 2022</a> )
<b>Fermented non-dairy products</b>				
<b>Sausages</b>	<i>Lactobacillus sakei</i> <i>Lactobacillus plantarum</i> , <i>Lactobacillus paracasei</i> subsp. <i>paracasei</i> , <i>Weissella viridescens</i> , <i>Lactobacillus coryniformis</i>	15 to 35°C for 1 to 5 days in the U.S for lab fermentation 20 to 32°C for 2 to 5 days in Europe pH for lab fermentation in Sausages is typically below 4.7	LAB present at end of sausages is about 5- 7 log CFU/g of sausages	( <a href="#">Leroy &amp; De Vuyst, 1999</a> ; <a href="#">Zamfir et al., 2022</a> ) ( <a href="#">Mohammadpourfard et al., 2021</a> ) ( <a href="#">Messens et al., 2003</a> )

LAB plays a crucial role in the food industry, particularly in cheese production. These bacteria are extensively used for various purposes in cheese making, contributing to the fermentation, flavor development, and quality of cheeses. In the food industry, LAB is majorly employed as starter cultures in cheese production, aiding in curd acidification and influencing the appearance and flavor of the end product. Additionally, LAB is utilized as a non-starter LAB (NSLAB) in cheese ripening, further enhancing the sensory characteristics of cheeses (Bettera et al., 2023; Valdiviezo-Marcelo et al., 2023). In previous studies, 16S rRNA genome sequencing plays an important role in studying the genome of ripening cheese, which is carried out by LAB strains. These studies give us the microbiota synthesis of smear-aged cheddar (cheese), and help us decide how individual bacterial species and microbial communities change over the long run inside the cheese grid. The microbiota progression was checked during the industrial cheese production, the temperature of ripening and storing as shown in the table, which included maturation and starting maturing at 15 °C, and after the production of cheese, which included putting away the result in the market at 4 °C for an ensuing time of as long as 40 days (Korena et al., 2023; Lovayová et al., 2015). The uses of LAB in cheese production, which is held in the food industry, are multifaceted. LAB is essential for acidifying milk, initiating fermentation, and producing specific flavors and textures in cheeses. Moreover, these bacteria contribute to the safety and quality of cheeses, playing a vital role in the ripening process and overall maturation of the product. LAB also has probiotic potential, offering additional health benefits to consumers when present in cheeses (Ahansaz et al., 2023; Kongo, 2013). Overall, the industrial application of LAB in cheese production in the food industry is extensive and essential for ensuring the desired characteristics, flavors, and quality of cheeses. LAB are versatile microorganisms that significantly impact the cheese-making process, making them indispensable in the food industry. The significance of LAB in cheddar creation is because of the occurrence of starter cultures and Nonstarter LAB. Starter societies, fundamentally *Lc. Lactic* or *S. thermophilus* is liable for converting lactose into lactic acid at a controlled rate (Hosken et al., 2023; Kongo, 2013). In the cycle, it brings about a progressive reduction in pH, which essentially affects different parts of cheddar production and eventually determines the cheddar's structure and standard. During the beginning phases of many cheese maturing, *Lb. delbrueckii* or *Lb. helveticus* (Coelho et al., 2022; Nicosia et al., 2023) assume a basic part, separating proteins, utilizing lactose, creating fragrant mixtures, and giving substrates that can be additionally consumed by other microbial groups, for example, NSLAB. NSLAB primarily incorporates into the facultative heterofermentative *Lactobacillus* sort, trailed by *Pediococcus pentosaceus* (Coelho et al., 2022; Kongo, 2013). They can affect the cheddar taste and surface because of the development of mixtures from the catabolism of amino acids, for the most part, methionine, fragrant amino acids, and extended-chain amino acids, notwithstanding the amalgamation of EPS. Moreover, bacteriocins, hydrogen peroxide, diacetyl, and CO<sub>2</sub> are likewise created by NSLAB, acting as bio-preservatives and contributing to the cheese's safety (Table 7) (Coelho et al., 2022; Kongo, 2013).

**Table 7.** Demonstrate the bacterial isolates present in different milk products.

Name Of Products	Isolate Of Bacteria	Reference
Milk	<i>Bacillus</i>	(Huck et al., 2007)
	<i>Campylobacter</i>	(Msalya, 2017)
	<i>Clostridium</i>	(Kabuki et al., 2004)
	<i>Salmonella</i>	(Mdegela et al., 2004)
	<i>S. aureus</i>	(Bastam et al., 2021)
	<i>S. intermedius</i>	
	<i>S. epidermis</i>	
	<i>S. agalactiae</i>	
	<i>Listeria monocytogenes</i>	
	<i>Mycobacteria</i>	
Yogurt	<i>E. coli</i>	(Hervert et al., 2017)
	Enterobacteriaceae	(Canganella et al., 1999)
	<i>P. paucimobilis</i>	(Yang & Yoon, 2022)
	<i>E. cloacae</i>	
	<i>L. monocytogenes</i>	
Cheese	<i>Listeria monocytogenes</i>	(El Marrakchi et al., 1993)
	<i>Pseudomonas</i>	(Al-Gamal et al., 2019)
	<i>Salmonella</i>	(Costanzo et al., 2020)
	<i>Bacillus</i>	
	<i>S. aureus</i>	



### 6.1.3. Food preservation

LAB assumes an urgent part in food safeguarding in the food business because of its capacity to hinder the development of a wide assortment of food decay microorganisms. They are known for their power to deliver restraining specialists, for example, lactic acid, which brings about a reduction of pH and improves the quality of food ([Negm, 2018](#); [Soomro, 2002](#)). LAB is added to food in society and is, by and large, viewed as innocuous or even helpful for human wellbeing in the US, where they are recorded as GRAS (For the most part Viewed as Safe). LAB produces antimicrobial mixtures called bacteriocins, which stand out enough to be noticed because of their possible handiness as regular substitutes for synthetic food additives. Bacteriocins-conveying social orders have been precisely applied to thwart *Listeria monocytogenes* and *Clostridium* spp. in various developed types of meat, vacuum-packaged products, and vegetable-based food assortments ([Mokoena et al., 2021](#)). Nisin A is one of the most important bacteriocins used as a food preservative. The spread of immunizing agent poison hindrance and interest in food things with fewer added substances requires a search for new decisions to avoid the abuse of accommodating serum poisons.

LAB withdrawn from uniquely designed developed vegetables produces antibacterial substances against both Gram-positive and Gram-negative ordinary foodborne bacterial microorganisms ([Mokoena et al., 2021](#); [Negm, 2018](#)). This wide variety of restrictions suggests that the LAB strains have the potential to be used as expected bio-preservatives in various food things, and to fight foodborne microorganisms. Bacteriocins are generally bactericidal, while some are bacteriostatic, conveying support in the food and medication regions, especially where development is poor. Bacteriocins are effective against Gram-positive toxigenic and pathogenic organisms, acting by forming pores in the membranes of target microorganisms. Heterofermentative *Lactobacillus* spp. have been shown to keep out wasted organic entities in cheddar processing. Bacteriocins can restrain the expansion of food decay microorganisms and foodborne microbes, making them a promising option in contrast to chemical additives in the dairy business ([Mezaini & Bouras, 2013](#); [Negm, 2018](#)).

## 7. Conclusion

LAB and probiotics are a non-chemical strategy for reducing food contamination through several mechanisms of action. The essential, organically important acids, bacteriocins, and other inhibitory compounds that LAB and probiotics produce, and their competition with harmful microorganisms, distinguish them from other existing cultures and position them as an essential part of food safety strategies. Because, as for the above, the successful application of LAB and probiotics in many food products, from dairy and meats to fermented vegetables and beverages, underlines their benefits: enhancing the quality, shelf life, and prevention of disease in foods. Increased consumer demand for natural, safe preservation requires deeper integration of LABs and probiotics in the food chain. Importantly, there is a need to investigate the best uses of these additives in different food systems and how they interact with various contaminants. Ultimately, this can sharply enhance the benefits of using LABs and probiotics in creating a much safer, sustainable, and healthy food supply chain.

### CRedit authorship contribution statement

We, the authors, declare that the manuscript titled "Role of *Lactobacillus* probiotics in gut health and food " Safety: Mechanisms for contaminant reduction and applications in food products" is an original work and has not been submitted elsewhere for publication. The authors – Mohsin Gulzar Barq, Amna Razzaq, Manal Yasin, Khadija Nawal, and Tayyaba Qureshi have all contributed significantly to the preparation of this review.

### Funding

No external funding was received for the preparation of this manuscript.

### Declaration of Competing Interest

The authors declared no conflict of interest.

### Acknowledgments

The authors gratefully acknowledge the valuable support in terms data analysis and technical assistance provided by Mr. Muhammad Hashir from the University of Central Punjab, Lahore, Pakistan.

## References

- Ağagündüz, D., Yılmaz, B., Şahin, T. Ö., Güneşliol, B. E., Ayten, Ş., Russo, P., Spano, G., Rocha, J. M., Bartkiene, E., & Özogul, F. (2021). Dairy lactic acid bacteria and their potential function in dietetics: The food-gut-health axis. *Foods*, 10(12), 3099.
- Ahansaz, N., Tarrah, A., Pakroo, S., Corich, V., & Giacomini, A. (2023). Lactic acid bacteria in dairy foods: Prime sources of antimicrobial compounds. *Fermentation*, 9(11), 964.
- Al-Gamal, M. S., Ibrahim, G. A., Sharaf, O. M., Radwan, A. A., Dabiza, N. M., Youssef, A. M., & El-Ssayad, M. F. (2019). The protective potential of selected lactic acid bacteria against the most common contaminants in various types of cheese in Egypt. *Heliyon*, 5(3).
- Almeida, K. E. d., Tamime, A. Y., & Oliveira, M. N. d. (2009). Influence of total solids contents of milk whey on the acidifying profile and viability of various lactic acid bacteria. *LWT-Food Science and Technology*, 42(2), 672-678.
- Arslan, S. (2015). A review: chemical, microbiological and nutritional characteristics of kefir. *CyTA-Journal of Food*, 13(3), 340-345.
- Bastam, M. M., Jalili, M., Pakzad, I., Maleki, A., & Ghafourian, S. (2021). Pathogenic bacteria in cheese, raw and pasteurised milk. *Veterinary Medicine and Science*, 7(6), 2445-2449.
- Bayat, M., Hashemi, J., & Razavilar, V. (2021). Identification of dairy fungal contamination and reduction of aflatoxin M1 amount by three acid and bile resistant probiotic bacteria. *Archives of Razi Institute*, 76(1), 119.
- Béjaoui, A., Ben Abdallah, I., & Maaroufi, A. (2022). *Brucella* spp. contamination in artisanal unpasteurized dairy products: An emerging foodborne threat in Tunisia. *Foods*, 11(15), 2269.
- Bettera, L., Levante, A., Bancalari, E., Bottari, B., & Gatti, M. (2023). Lactic acid bacteria in cow raw milk for cheese production: Which and how many? *Frontiers in microbiology*, 13, 1092224.
- Bintsis, T. (2018). Lactic acid bacteria: their applications in foods. *J. Bacteriol. Mycol*, 6(2), 89-94.
- Buehler, A., Martin, N., Boor, K., & Wiedmann, M. (2018). Evaluation of biopreservatives in Greek yogurt to inhibit yeast and mold spoilage and development of a yogurt spoilage predictive model. *Journal of dairy science*, 101(12), 10759-10774.
- Butler, M. I., Bastiaanssen, T. F., Long-Smith, C., Berding, K., Morkl, S., Cusack, A.-M., Strain, C., Busca, K., Porteous-Allen, P., & Claesson, M. J. (2020). Recipe for a healthy gut: intake of unpasteurised milk is associated with increased lactobacillus abundance in the human gut microbiome. *Nutrients*, 12(5), 1468.
- Cálix-Lara, T. F., Rajendran, M., Talcott, S. T., Smith, S. B., Miller, R. K., Castillo, A., Sturino, J. M., & Taylor, T. M. (2014). Inhibition of *Escherichia coli* O157: H7 and *Salmonella enterica* on spinach and identification of antimicrobial substances produced by a commercial Lactic Acid Bacteria food safety intervention. *Food Microbiology*, 38, 192-200.
- Canganella, F., Nespica, M., Giontella, D., & Trovatelli, L. (1999). Survival of *Enterobacter cloacae* and *Pseudomonas paucimobilis* in yoghurts manufactured from cow's milk and soymilk during storage at different temperatures. *Microbiological research*, 154(1), 15-21.
- Chen, C., Zhao, S., Hao, G., Yu, H., Tian, H., & Zhao, G. (2017). Role of lactic acid bacteria on the yogurt flavour: A review. *International Journal of Food Properties*, 20(sup1), S316-S330.
- Coelho, M. C., Malcata, F. X., & Silva, C. C. (2022). Lactic acid bacteria in raw-milk cheeses: From starter cultures to probiotic functions. *Foods*, 11(15), 2276.
- Collado, M., Meriluoto, J., & Salminen, S. (2007). Role of commercial probiotic strains against human pathogen adhesion to intestinal mucus. *Letters in applied microbiology*, 45(4), 454-460.
- Colombo, M., Castilho, N. P., Todorov, S. D., & Nero, L. A. (2018). Beneficial properties of lactic acid bacteria naturally present in dairy production. *BMC microbiology*, 18(1), 219.
- Costanzo, N., Ceniti, C., Santoro, A., Clausi, M. T., & Casalnuovo, F. (2020). Foodborne pathogen assessment in raw milk cheeses. *International journal of food science*, 2020(1), 3616713.
- da Silva, T. M. S., Piazentin, A. C. M., Mendonça, C. M. N., Converti, A., Bogsan, C. S. B., Mora, D., & de Souza Oliveira, R. P. (2020). Buffalo milk increases viability and resistance of probiotic bacteria in dairy beverages under in vitro simulated gastrointestinal conditions. *Journal of dairy science*, 103(9), 7890-7897.
- Dehkordi, F. S., Yazdani, F., Mozafari, J., & Valizadeh, Y. (2014). Virulence factors, serogroups and antimicrobial resistance properties of *Escherichia coli* strains in fermented dairy products. *BMC research notes*, 7(1), 217.
- Di Stefano, V. (2014). Food contaminants. *Journal of Food Studies*.



- Ei-Shafe, H. M., Abo-Zaid, K. F., & Madkour, A. M. (2017). Effect Of Microbial Contamination With Fungi, Aflatoxin M1 And Enteric Gram Negative Bacteria On Milk And Some Dairy Products. *Egyptian Journal of Agricultural Research*, 95(4), 1871-1889.
- El Marrakchi, A., Hamama, A., & El Othmani, F. (1993). Occurrence of *Listeria monocytogenes* in milk and dairy products produced or imported into Morocco. *Journal of Food Protection*, 56(3), 256-259.
- Gänzle, M. G., Monnin, L., Zheng, J., Zhang, L., Coton, M., Sicard, D., & Walter, J. (2024). Starter culture development and innovation for novel fermented foods. *Annual review of food science and technology*, 15.
- Hervert, C., Martin, N., Boor, K., & Wiedmann, M. (2017). Survival and detection of coliforms, Enterobacteriaceae, and gram-negative bacteria in Greek yogurt. *Journal of dairy science*, 100(2), 950-960.
- Hojsak, I., Abdović, S., Szajewska, H., Milošević, M., Krznarić, Ž., & Kolaček, S. (2010). *Lactobacillus* GG in the prevention of nosocomial gastrointestinal and respiratory tract infections. *Pediatrics*, 125(5), e1171-e1177.
- Hosken, B. d. O., Melo Pereira, G. V., Lima, T. T. M., Ribeiro, J. B., Magalhães Júnior, W. C. P. d., & Martin, J. G. P. (2023). Underexplored potential of lactic acid bacteria associated with artisanal cheese making in Brazil: Challenges and opportunities. *Fermentation*, 9(5), 409.
- Huck, J., Hammond, B., Murphy, S., Woodcock, N., & Boor, K. (2007). Tracking spore-forming bacterial contaminants in fluid milk-processing systems. *Journal of dairy science*, 90(10), 4872-4883.
- Hussain, M. A., & Gooneratne, R. (2017). Understanding the fresh produce safety challenges. In (Vol. 6, pp. 23): MDPI.
- Ibrahim, S. A., Ayivi, R. D., Zimmerman, T., Siddiqui, S. A., Altemimi, A. B., Fidan, H., Esatbeyoglu, T., & Bakhshayesh, R. V. (2021). Lactic acid bacteria as antimicrobial agents: Food safety and microbial food spoilage prevention. *Foods*, 10(12), 3131.
- Ismail, M., & Sabreen, M. (2001). Associated Mycobiota Of Some Types Of Cheese And Cooking Butter. *Assiut Veterinary Medical Journal*, 44(88), 176-197.
- Kabuki, D., Kuaye, A., Wiedmann, M., & Boor, K. (2004). Molecular subtyping and tracking of *Listeria monocytogenes* in Latin-style fresh-cheese processing plants. *Journal of dairy science*, 87(9), 2803-2812.
- Khatoun, S., Kalam, N., Rashid, S., & Bano, G. (2023). Effects of gut microbiota on neurodegenerative diseases. *Frontiers in Aging Neuroscience*, 15, 1145241.
- Kim, H. S., & Gilliland, S. E. (1983). *Lactobacillus acidophilus* as a dietary adjunct for milk to aid lactose digestion in humans. *Journal of dairy science*, 66(5), 959-966.
- Kongo, J. M. (2013). *Lactic Acid Bacteria: R. BoD-Books on Demand*.
- Korena, K., Krzyzankova, M., Florianova, M., Karasova, D., Babak, V., Strakova, N., & Juricova, H. (2023). Microbial succession in the cheese ripening process—Competition of the starter cultures and the microbiota of the cheese plant environment. *Microorganisms*, 11(7), 1735.
- Leroy, F. d. r., & De Vuyst, L. (1999). Temperature and pH conditions that prevail during fermentation of sausages are optimal for production of the antilisterial bacteriocin sakacin K. *Applied and environmental microbiology*, 65(3), 974-981.
- López, P., & Spano, G. (2023). Industrial and health applications of lactic acid bacteria and their metabolites, volume II. In (Vol. 14, pp. 1242253): Frontiers Media SA.
- Lovayová, V., Dudriková, E., Rimárová, K., & Siegfried, L. (2015). Quantity of selected probiotic cultures in semi-hard cheese with low-cooking curd during the maturation process. *Journal of food science and technology*, 52(8), 4697-4702.
- Maldonado, J., Lara-Villoslada, F., Sierra, S., Sempere, L., Gomez, M., Rodriguez, J. M., Boza, J., Xaus, J., & Olivares, M. (2010). Safety and tolerance of the human milk probiotic strain *Lactobacillus salivarius* CECT5713 in 6-month-old children. *Nutrition*, 26(11-12), 1082-1087.
- Mantzourani, I., Nouska, C., Terpou, A., Alexopoulos, A., Bezirtzoglou, E., Panayiotidis, M. I., Galanis, A., & Plessas, S. (2018). Production of a novel functional fruit beverage consisting of cornelian cherry juice and probiotic bacteria. *Antioxidants*, 7(11), 163.
- Marcos, A., Wörnberg, J., Nova, E., Gómez, S., Alvarez, A., Alvarez, R., Mateos, J. A., & Cobo, J. M. (2004). The effect of milk fermented by yogurt cultures plus *Lactobacillus casei* DN-114001 on the immune response of subjects under academic examination stress. *European journal of nutrition*, 43(6), 381-389.
- Mazahreh, A. S., & Ershidat, O. T. M. (2009). The benefits of lactic acid bacteria in yogurt on the gastrointestinal function and health. *Pakistan Journal of Nutrition*, 8(9), 1404-1410.
- Mdegela, R., Kusiluka, L., Kapaga, A., Karimuribo, E., Turuka, F., Bundala, A., Kivaria, F., Kabula, B., Manjurano, A., & Loken, T. (2004). Prevalence and determinants of mastitis and milk-borne zoonoses

- in smallholder dairy farming sector in Kibaha and Morogoro districts in Eastern Tanzania. *Journal of Veterinary Medicine, Series B*, 51(3), 123-128.
- Messens, W., Verluyten, J., Leroy, F., & De Vuyst, L. (2003). Modelling growth and bacteriocin production by *Lactobacillus curvatus* LTH 1174 in response to temperature and pH values used for European sausage fermentation processes. *International journal of food microbiology*, 81(1), 41-52.
- Mezaini, A., & Bouras, A. D. (2013). Antibacterial activity and probiotic properties of some lactic acid bacteria isolated from dairy products. *African Journal of Biotechnology*, 12(20).
- Miranda, C., Contente, D., Igrejas, G., Câmara, S. P., Dapkevicius, M. d. L. E., & Poeta, P. (2021). Role of exposure to lactic acid bacteria from foods of animal origin in human health. *Foods*, 10(9), 2092.
- Mohammadpourfard, I., Khanjari, A., Akhonzadeh Basti, A., Herrero-Latorre, C., Shariatifar, N., & Hosseini, H. (2021). Evaluation of microbiological, chemical, and sensory properties of cooked probiotic sausages containing different concentrations of astaxanthin, thymol, and nitrite. *Food science & nutrition*, 9(1), 345-356.
- Mokoena, M. P., Omatola, C. A., & Olaniran, A. O. (2021). Applications of lactic acid bacteria and their bacteriocins against food spoilage microorganisms and foodborne pathogens. *Molecules*, 26(22), 7055.
- Msalya, G. (2017). Contamination levels and identification of bacteria in milk sampled from three regions of Tanzania: evidence from literature and laboratory analyses. *Veterinary Medicine International*, 2017(1), 9096149.
- Negm, G. E. S. (2018). Classification, antimicrobial potential, industrial applications and probiotic capability of lactic acid bacteria: A review article. *Research Journal of Applied Sciences*, 13(12), 742-757.
- Nicosia, F. D., Pino, A., Maciel, G. L. R., Sanfilippo, R. R., Caggia, C., de Carvalho, A. F., & Randazzo, C. L. (2023). Technological characterization of lactic acid bacteria strains for potential use in cheese manufacture. *Foods*, 12(6), 1154.
- Omar, D., Al-Ashmawy, M., Ramadan, H., & El-Sherbiny, M. (2018). Occurrence and PCR identification of *Salmonella* spp. from milk and dairy products in Mansoura, Egypt. *International Food Research Journal*, 25(1), 446-452.
- Onoharigho, F. O., Ighede, P. A., Edo, G. I., Akpoghelie, P. O., & Akpoghelie, E. O. (2022). Isolation and identification of bacterial and fungal spoilage organisms in branded and unbranded milk; consumer perception of safety hazard for milk. *Applied Microbiology: Theory & Technology*, 31-48.
- Parada, J. L., Caron, C. R., Medeiros, A. B. P., & Soccol, C. R. (2007). Bacteriocins from lactic acid bacteria: purification, properties and use as biopreservatives. *Brazilian Archives of Biology and Technology*, 50(3), 512-542.
- Pei, X., Tekliye, M., & Dong, M. (2021). Isolation and identification of fungi found in contaminated fermented milk and antifungal activity of vanillin. *Food Science and Human Wellness*, 10(2), 214-220.
- Pereg, D., Kimhi, O., Tirosh, A., Orr, N., Kayouf, R., & Lishner, M. (2005). The effect of fermented yogurt on the prevention of diarrhea in a healthy adult population. *American journal of infection control*, 33(2), 122-125.
- Pérez-Rivero, C., & López-Gómez, J. P. (2023). Unlocking the potential of fermentation in cosmetics: A review. *Fermentation*, 9(5), 463.
- Prado, M. R., Blandón, L. M., Vandenberghe, L. P., Rodrigues, C., Castro, G. R., Thomaz-Soccol, V., & Soccol, C. R. (2015). Milk kefir: composition, microbial cultures, biological activities, and related products. *Frontiers in microbiology*, 6, 1177.
- Rather, I. A., Koh, W. Y., Paek, W. K., & Lim, J. (2017). The sources of chemical contaminants in food and their health implications. *Frontiers in pharmacology*, 8, 830.
- Rostami, F., Rahimi, E., Yahaghi, E., Khodaverdi Darian, E., & Bagheri Moghadam, M. (2014). Isolation and evaluation virulence factors of *Salmonella typhimurium* and *Salmonella enteritidis* in milk and dairy products. *Iranian Journal of Medical Microbiology*, 8(1), 54-61.
- Serpe, C. J., Kohm, A. P., Huppenbauer, C. B., Sanders, V. M., & Jones, K. J. (1999). Exacerbation of facial motoneuron loss after facial nerve transection in severe combined immunodeficient (scid) mice. *The Journal of Neuroscience*, 19(11), RC7.
- Soomro, A. (2002). Role of lactic acid bacteria (LAB) in food preservation and human health a review. *Pak. J. Nutr.*
- Tan, L. L., Tan, C. H., Ng, N. K. J., Tan, Y. H., Conway, P. L., & Loo, S. C. J. (2022). Potential probiotic strains from milk and water kefir grains in Singapore – use for defense against enteric bacterial pathogens. *Frontiers in microbiology*, 13, 857720.



- Valdiviezo-Marcelo, J., Arana-Torres, N. M., Vega-Portalatino, E. J., Ruiz-Flores, L. A., Tamariz-Angeles, C., Olivera-Gonzales, P., Rosales-Cuentas, M. M., & Espinoza-Espinoza, L. A. (2023). Technological potential of native lactic acid bacteria isolated from Swiss-type artisanal cheese (Ancash, Peru) for their application in food. *Frontiers in Sustainable Food Systems*, 7, 1212229.
- Wang, Y., Wu, J., Lv, M., Shao, Z., Hungwe, M., Wang, J., Bai, X., Xie, J., Wang, Y., & Geng, W. (2021). Metabolism characteristics of lactic acid bacteria and the expanding applications in food industry. *Frontiers in bioengineering and biotechnology*, 9, 612285.
- Worku, K. F., Kurabachew, H., & Hassen, Y. (2019). Probiotication of Fruit Juices by Supplemented Culture of lactobacillus acidophilus. *Int. J. Food Sci. Nutr. Eng*, 9, 45-48.
- Yang, S. Y., & Yoon, K. S. (2022). Quantitative microbial risk assessment of *Listeria monocytogenes* and enterohemorrhagic *Escherichia coli* in yogurt. *Foods*, 11(7), 971.
- Zamblé, B., Lessoy, Z., & Rose, K.-N. (2016). FUNGAL CONTAMINATION AND MYCOTOXINS' OCCURRENCE IN PEANUT BUTTERS MARKETED IN ABIDJAN DISTRICT (CÔTE D'IVOIRE). *Food and Environment Safety Journal*, 13(3).
- Zamfir, M., Angelescu, I.-R., Voaides, C., Cornea, C.-P., Boiu-Siculia, O., & Grosu-Tudor, S.-S. (2022). Non-dairy fermented beverages produced with functional lactic acid bacteria. *Microorganisms*, 10(12), 2314.
- Zapašnik, A., Sokołowska, B., & Bryła, M. (2022). Role of lactic acid bacteria in food preservation and safety. *Foods*, 11(9), 1283.
- Zhang, Z., Lv, J., Pan, L., & Zhang, Y. (2018). Roles and applications of probiotic *Lactobacillus* strains. *Applied microbiology and biotechnology*, 102(19), 8135-8143.