



Characterization Of Bioactive Compounds In Luffa Aegyptiaca Leaf Ethanolic Extracts Using Gas Chromatography And Mass Spectrometer

Alagbe, J.O^{a*}, Bamigboye Samson^b, G.C. Nwosu^c, D.A Agbonika^d, Kadiri Mercy Cincinsoko^e

a. Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India

b, c. Department of Animal Science, University of Abuja, Nigeria

d. Department of Agricultural Economics, University of Abuja, Nigeria

e. Kazachat College Abuja, Nigeria.

Abstract: Medicinal plants contain several bioactive compounds or phytochemicals that therapeutically exert wide spectrum of biological activities such as hepato-protective, hypolipidemic, analgesics, antioxidant, immune-stimulatory, immune-modulatory and anticancer. These bioactive compounds are effective, non-toxic and environmentally friendly. Here we determined the bioactive compounds in Luffa aegyptiaca ethanolic leaf extract using gas chromatography and mass spectrometry (GC-MS). Biochemical analysis revealed the presence of alkaloids (1314.9 mg/g), glycosides (11.60 mg/g) while tannins, flavonoids, terpenoids, steroids, saponins and oxalates were present in 807.56 mg/g, 512.9 mg/g, 308.7 mg/g, 192.6 mg/g, 102.1 mg/g, 61.44 mg/g and 20.90 mg/g respectively. GC-MS analysis of Luffa aegyptiaca ethanolic leaf extract shows that it contains 35 bioactive compounds with marked pharmaceutical properties. 1,11-Bicyclopropyl] 2-octanoic acid had the highest concentration (19.06 %) while β -methylnaphthalene had the lowest concentration (0.01 %). It was concluded that the presence of these phytochemicals will inhibit some pathogenic bacteria's thus promoting a healthy gut in animals.

Keywords: Luffa aegyptiaca extracts, medicinal plants, phytochemicals, gas chromatography, mass spectrometry

1. Introduction:

Medicinal plants have been generally recognized to contain phytochemicals or secondary metabolites which performs multiple physiological functions (Alagbe, 2023; Singh et al., 2022). They have globally gained interest due to their efficacy, safety and environmental friendly with lesser side effects (Adewale et al., 2021; Singh et al., 2021). In developing countries, eighty percent of world population use medicinal plants for primary health care and it has been estimated that nearly 75 % of plants contain several bioactive compounds of therapeutic properties (Shittu et al., 2022; Alagbe, 2021).

Luffa aegyptiaca is one of the most potent herbs with long history of medicinal use. The plant belongs to the family Cucurbitaceae and order Cucurbitales (Alagbe, 2019; Farag et al., 2015). It is also known as sponge gourd and widely grown in Asia, Europe and many parts of Africa (Parkash and Tso, 2002; Muthumani et al., 2010). Various parts of the plant (seeds, leaves, root and stem bark) have been documented as a traditional treatment for fever, gastro intestinal infections, convulsion, skin diseases, diabetes, rheumatism, snake bite and back ache (Irshad et al., 2010; Azeez et al., 2013). It can also display anti-inflammatory (Kao et al., 2012), immuno-modulatory, immune-stimulants, antiviral (Ng et al., 2011), antifungal, hypoglycemic, cytotoxic and antioxidant properties due to the presence of some secondary metabolites (Lawal et al., 2010; Farag et al., 2012; Smith et al., 2006).

Infused methanolic and aqueous leaf extract from Luffa aegyptiaca leaves are capable of inhibiting the growth of pathogenic bacteria's (Gram -ve and Gram +ve) (Roy and Lingampeta, 2014; Wildman, 2001). Isolates from the seeds and aqueous stem bark extract (Luffacylin and dihydroxy spinasterol) have been reported to exhibit

[Received 09 Sep 2022; Accepted 17 Nov 2022; Published (online) 22 Dec 2022]

Finesse Publishing stays neutral regard to jurisdictional claims published maps



Attribution 4.0 International (CC BY 4.0)

Corresponding email: dralagbe@outlook.com (Alagbe, J.O)

DOI: 10.61363/fsamr.v1i2.53

antifungal and antimicrobial activities due to the presence of pharmacologically active substances (Koledoye et al., 2021; Amin et al., 2009). Characterization of *Luffa aegyptiaca* leaves to ascertain their phytoconstituents using gas chromatography and mass spectrometry will help to develop novel drugs and its efficacy against wider range of pathogenic organisms (Patel et al., 2013; Ediriweera et al., 2019). Here we determine the bioactive compounds in *Luffa aegyptiaca* leaf extract.

2. Materials and methods

2.1 Experimental site, plant collection, processing and phytochemical analysis

Fresh *Luffa aegyptiaca* leaves were collected from Punsari village, Gujarat and transferred to Sumitra Research Institute where it was authenticated by a certified taxonomist. Dried leaves were blended into powder using an electric blender to allow easy penetration of solvent (ethanol). 200 grams of the blended sample was transferred into Erlenmeyer's flask followed by the addition of 1000 liters of 95 % ethyl alcohol, stirred occasionally at intervals for 24 hours and filtered into another flask using Whatman's filter paper thereafter it was covered with aluminum foil to avoid the solvent from escaping the mixture. It was later set on a water bath at 70 oC for 10 minutes to recover the extract for further laboratory examination.

Quantitative phytochemical analysis were carried out using standard laboratory procedures Tannins was analyzed using methods outlined by Biswas et al. (2020), flavonoids and terpenoids (Surana et al., 2016), phenols, alkaloids and saponins (Madhu et al., 2016), phytates and oxalates (He et al., 2014) and steroids (Adeniyi et al., 2017).

2.2 Analysis of secondary metabolites in *Luffa aegyptiaca* leaves using GC-MS

Determination of secondary metabolites in ethanolic *Luffa aegyptiaca* leaf extract was done with LABTRON gas chromatography mass spectrometry (GC-MS-879) by following standard protocol.

Table 1: secondary metabolites in *Luffa aegyptiaca* leaves using Gas chromatography and mass spectrometry

Compounds	Reaction time (Sec)	% Area
2-methoxy-4-vinylphenol	3.561	0.95
β -elemenone	4.700	5.67
Erythritol	4.766	8.40
7-methylenebicyclo hepata -3-ene	5.006	3.81
Guanosine	5.882	8.56
α -terpineol	6.881	0.18
Limonene	7.009	1.25
Isorbide dinitrate	8.012	0.09
Heptadec-3-enal	8.116	2.10
Ethyl Oleate	9.301	0.04
9,15-Octadecadienoic acid	10.04	0.06
Dibutyl benzene-1,2 dicarboxylate	10.33	1.85



Diethyl suberate	11.67	0.03
4-Acetoxy-3-methoysterene	11.88	0.46
Camphol	16.09	5.03
β -Linalool	20.06	3.60
6,6-Dimethyl -1,3-heptadien-5-ol	24.09	1.25
Benzaldehyde	28.00	0.73
β -methylnaphthalene	28.82	0.01
3-Octanone	29.04	0.55
Isomycorene	30.03	0.08
Dihydromyrcenol	31.40	1.45
Glucopyranoside	32.67	4.03
Spirost-8-en-11-one	35.09	1.22
2-Nonenoic acid	36.02	1.78
Ethyl iso-allocholate	37.01	0.79
Monomethyl pimelate	38.30	2.06
4-Acetoxy-3-methoxystyrene	40.10	0.02
2,4,6 – Octarien -1-ol 3,7 dimethyl	42.08	0.05
Lutoelin	45.66	0.03
1,11-Bicyclopropyl] 2-octanoic acid	46.08	19.06
Apigenin	47.01	6.01
1-methylcyclopropanemethanol	47.66	0.02
Total		92.69

3. Results and discussion

3.1 Phytochemical composition of *Luffa aegptiaca* leaf extract

Phytochemicals or secondary metabolites are chemical compounds produced by plants and it possess medicinal (anti-inflammatory, antidiuretic, immune-stimulant, anticancer, immune-modulatory, cytotoxic, hepato protective, antiviral etc.) and also have many applications (pharmaceutical, agrochemicals and cosmetics) (Shittu and Alagbe, 2021). Phytochemical composition of *Luffa aegptiaca* leaf extract shows that Alkaloids had the highest concentration (1314.9 mg/g) while glycosides had the lowest concentration (11.60 mg/g). Tannins, flavonoids, terpenoids, oxalates, saponins steroids and were present in 807.56 mg/g, 512.9 mg/g, 308.7 mg/g, 192.6 mg/g, 102.1 mg/g, 61.44 mg/g and 20.90 mg/g respectively. The higher concentration in alkaloid values is a clear sign that the extract can invoke a bitter taste and can perform wide range of pharmacological activities (Alagbe, 2021; Shittu et al., 2021; Agubosi et al., 2022). Glycosides are organic compounds made up of sugars and aglycone. It can exist in various forms (cardiac glycosides, hydroquinone glycosides and anthraquinone glycosides) and has antiseptic, expectorant, antihyperlipidemic and rubefacient (Parama et al., 2020; Elekofehinti et al., 2015). Flavonoids are group of phenolic compounds with antibacterial, antiviral effects and prevention of deoxyribonucleic acid binding (Alagbe et al., 2020; Alagbe and Omokore, 2019). It also aids in enzyme induction and enhancing detoxification of organs (Shmuel, 2012). Terpenoids therapeutically exerts wide spectrum of properties such as; antibacterial, anti-helminthic, antioxidant, analgesics, hypolipidemic and cytotoxic functions (Park et al., 2012; Alagbe, 2019). Tannins are naturally occurring complex compounds which possess nitrogen free polyphenols with several pharmaceutical effects (antimicrobial, cytotoxic, antioxidant and hepato-protective properties) (Tanaka et al., 1991). High concentration of oxalate in the system of animals could result in kidney stone disease and osteoporosis (Griego et al., 2008).




Figure 1: Phytochemical composition of *Luffa aegptiaca* leaf extract

4. Secondary metabolites in *Luffa aegptiaca* leaf extract using gas chromatography and mass spectrometry

The result reveals the presence of 35 bioactive compounds that are capable of giving therapeutic effect on the health of human and animals (Urbanova et al., 2013; Fan and Qian, 2006). These compounds includes: 2-methoxy-4-vinylphenol (0.95 %), β -elemenone (5.67 %), Erythritol (8.40 %), 7-methylenebicyclo hepata -3-ene (3.81 %), Guanosine (8.56 %), α -Terpineol (0.18 %), Limonene (1.25 %), Isorbide dinitrate (0.09 %), Heptadec-3-enal (2.10 %), Ethyl Oleate (0.04 %), 9,15-Octadecadienoic acid (0.06 %), Dibutyl benzene-1,2 dicarboxylate (1.85 %), Diethyl suberate (0.03 %), 4-Acetoxy-3-methoysterene (0.46 %), Camphol (5.03 %), β -Linalool (3.60 %), 6,6-Dimethyl -1,3-heptadien-5-ol (1.25 %), Benzaldehyde (0.73 %), β -methylnaphthalene (0.01 %), 3-Octanone (0.55 %), Isomycorene (0.08 %), Dihydromyrcenol (1.45 %), Glucopyranoside (4.03 %), Estra-1,3,5 (10) -trien-17-ol (0.57 %), Spirost-8-en-11-one (1.22 %), 2-Nonenoic acid (1.78 %), Ethyl iso-allocholate (0.79 %), Monomethyl pimelate (2.06 %), Hexan-3-yl 2-methylpropyl benzene 1,2 carboxylate (0.10 %), 4-Acetoxy-3-methoxystyrene (0.02 %), 2,4,6 - Octarien -1-ol 3,7 dimethyl (0.05 %), 2R, 3S-1-[1,3,4 - Trihydroxy-2-butoxymethyl (10.80 %), Lutoelin (0.03 %), 1,11-Bicyclopropyl] 2-octanoic acid (19.06 %), Apigenin (6.01 %) and 1-methylcyclopropanemethanol (0.02 %). These compounds play a vital role in traditional herbal



remedies and also have several pharmaceutical or physiological effect on human and animal (Zhang et al., 2006; Tang, 1992). The outcome on the GC-MS analysis of *Luffa aegyptiaca* leaf extract agrees with the reports of Garia et al. (2018); Yadav et al. (2017) but contrary to the findings of Akther et al. (2014). This disparity could be attributed to geographical location, species or parts of plant used (root, seed, stem bark, flower etc.), extraction methods, harvesting procedure and age of plants (Agubosi et al., 2022; Alagbe and Akintayo, 2020)..

5. Conclusion

The result on the phytochemical composition of *Luffa aegyptiaca* leaf extract confirms that it contains several bioactive compounds of marked pharmacological activities (antimicrobial, antifungal, antiviral, antioxidant, immune-stimulatory, hepato-protective, cytotoxic, hypolipidemic, immuno-modulatory, analgesics, and antihelminthic) and are also capable of inhibiting the growth of pathogenic bacteria.

6. Reference:

- Alagbe, J.O. (2023). Bioactive compounds in ethanolic extract of *Strychnos innocua* root using gas chromatography and mass spectrometry (GC-MS). *Drug Discovery*, 2023; 17:e4dd1005.
- Muritala, Daniel Shittu., Alagbe, J.O., Ojebiyi, O.O., Ojediran, T.K and Rafiu, T.A. (2022). Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of *Polyalthia longifolia* leaf extract in place of conventional antibiotics. *Animal Science and Genetics* 18(2): 57-71.
- Nascimento GGF, Lacatelli J, Freitas PC, Silva GL. (2012). Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. *Brazil Journal Microbiology*, 31(4):886-891.
- Okeke MI, Iroegbu CU, Eze EN, Okoli AS, Esimone C.O. (2001). Evaluation of extracts of the roots of *Landolphia owerience* for antibacterial activity. *Journal Ethnopharmacology*. 78(2-3):119-27.
- Jisika M, Ohigashi H, Nogaka H, Tada T, Hirota M. (1992). Bitter steroid glycosides, Vernonia sides A1, A2, and A3 and related B1 from the possible medicinal plant *vernonia amygdalina* used by wild Chimpanzees. *Tetrahedron*, 48:625-630.
- Jones GA, McAllister TA, Muir AD, Cheng K.J. (1996). Effects of safonin (*Onobrychis viciifolia scop.*) condensed tannins on growth and proteolysis by four strains of ruminal bacteria. *Appl. Environ. Microbiology*, 60:1374- 1378.
- Elumalai A, Chinna Eswaraiyah M. A. (2011). Pharmacological Review on *Garcinia indica.*, *International journal of universal pharmacy and Life Sciences*, 1(3):57-60.
- Narayani M, Johnson M, Sivaraman A, Janakiraman N. Phytochemical and Antibacterial Studies on *Jatropha curcas L.* *Journal of Chemical and Pharmaceutical Research*. 2012; 4(5):2639-2642.
- Parekh J, Chands S. In vitro antibacterial activity of the crude methanol extract of *Woodfordia fruticosa* Kurz. Flower (Lythraceae), *Brazil Journal of Microbiology*. 2007; 38:204-7.
- Adeyemi, A.O., Alagbe, J.O., Adeyemi, A.O. (2021). Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phytogenic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. *International Journal of Orange Technologies*, 3(3): 1-12.
- Alagbe, J.O and Akintayo-Balogun, O.M. (2020). Effects of dietary supplementation of *Albizia lebeck* seed oil (ALO) on the fatty acid composition of weaner rabbits. *Biochemistry and Biotechnology Research*, 8(2): 29-33.
- Akintayo Balogun Omolere. M and Alagbe, J.O (2020). Probiotics and medicinal plants in poultry nutrition: A review. *United International Journal for Research and Technology*, 2(1): 7-13.
- Alagbe, J.O. (2020). Effect of dietary supplementation of *Cymbopogon Citratus* oil on The Performance and Carcass characteristics of broiler chicks. *European Journal of Biotechnology and Bioscience*. 8(4): 39-45.
- Alagbe, J.O., Agubosi, O.C.P., Ajagbe, A.D, Shittu, M.D and Akintayo Balogun, O.M (2020). Performance, haematology and serum biochemical parameters of growing grass cutters fed *Phyllanthus amarus* and *Piliostigma thonningii* leaf meal mixture as partial replacement for Soya bean meal. *United International Journal for Research and Technology*, 2(1): 14-23.
- Alagbe, J.O., Shittu, M.D., Bamigboye, S.O and Oluwatobi, A.O. (2020). Proximate and mineral composition of *Pentadiplandra brazzeana* stems bark. *Electronic Research Journal of Engineering, Computer and Applied Sciences*. 1(2009): 91-99.

- Alagbe John Olujimi, Ramalan Sadiq Muhammad., Shittu Muritala Daniel and Olagoke Olayemi Christiana (2022). Effect of *Trichilia monadelpha* stem bark extract on the fatty acid composition of rabbit's thigh meat. *Journal of Environmental Issues and Climate Change* 1(1): 63-71.
- Alagbe, J.O., Shittu, M.D and Tanimomo, Babatunde K. (2022). Influence of *Anogeissus leio carpus* stem bark on the fatty acid composition in meat of broiler chickens. *European Journal of Life Safety and Stability* 14(22): 13-22.
- Alagbe, J.O (2022). Use of medicinal plants as a panacea to poultry production and food security: A review. *Gospodarka I Innowacje* 22(2022): 1-12.
- Singh Sharma., Alagbe Olujimi John., Liu Xing., Sharma Ram and Kumar Amita (2022). Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography and mass spectroemetry. *International Journal of Agriculture and Animal Production*, 2(6): 18-27.
- Agubosi, O.C.P., Alexander, James and Alagbe, J.O. (2022). Influence of dietary inclusion of Sunflower (*Helianthus annuus*) oil on growth performance and oxidative status of broiler chicks. *Central Asian Journal of Medical and Natural Sciences* 2(7): 187-195.
- Agubosi, O.C.P., Soliu, M.B and Alagbe, J.O. (2022). Effect of dietary inclusion levels of *Moringa oleifera* oil on the growth performance and nutrient retention of broiler starter chicks. *Central Asian Journal of Theoretical and Applied Sciences* 3(3): 30-39.
- Alagbe, J.O. (2022). Gas chromatography and mass spectroscopy of *Juniperus phoenice* stem bark extract and its influence on the haemato-biochemical values of growing rabbits. *British Scientific Periodical* 1(1): 18-33.
- Alagbe, J.O., Zubairu Habiba., Adedeji, O.M., Bamigboye, S and Dora Agbonika (2022). Influence of *Juniperus thurifera* root extract on the nutrient digestibility and caecal microbial count of growing rabbits. *Web of Synergy: International Interdisciplinary Research Journal* 1(1): 5-17.
- Alagbe, J.O., Shittu, M.D., Ramalan, S.N., Tanimomo, K.B and Adekunle, D.A. (2022). Growth performance, semen quality characteristics and hormonal profile of male rabbit bucks fed *Rubia cordifolia* root extracts. *International Journal of Biological Engineering and Agriculture* 1(1): 1-13.
- Agubosi, O.C.P., Imudia, Favour Dumkenechukwu and Alagbe, J.O. (2022). Evaluation of the nutritional value of air dried and sun-dried sweet potato (*Ipomoea batatas*) peels. *European Journal of Life Safety and Stability* 14(22): 43-51.
- Roy S. and Lingampeta P. (2014). Solid wastes of fruits peels as source of low cost broad spectrum natural antimicrobial compounds furanone, furfural and benzenetriol. *International Journal of Research in Engineering and Technology*, 3(7): 273-279.
- Wildman, R.E.C. (2001). Classifying nutraceuticals. In *Handbook of nutraceuticals and functional foods* (Edn. Wolinsky, Hickson and J.F. Jr., I). CRC press LLC.
- Hung, P.V., Maeda, T., Miyatake, K. and Morita, N. Total Phenolic Compounds and Antioxidant Capacity of Wheat Graded Flours by Polishing Method. *Food Res Int.* 42(1) 185-190 (2009).
- Lawal I, Uzokwe N, Igboanugo A, Adio A, Awosan E, Nwogwugwu J. (2010). Ethno medicinal information on collation and identification of some medicinal plants in Research Institutes of South-west Nigeria. *African Journal of Pharmaceutical Pharmacology*, 4 (1):001-7.
- Azeez MA, Bello OS, Adedeji A.O. (2013). Traditional and medicinal uses of *Luffa cylindrica*: a review. *Journal of Medicinal Plants*. 1(5):102-11.
- Irshad IA M, Goel HC, Moshahid M, Rizvi A. (2010). Phytochemical screening and high performance TLC analysis of some cucurbits. *Res J Phytochemicals*, 4 (4):242-7
- Parkash A, Ng T, Tso W. (2002). Isolation and characterization of luffacylin, a ribosome inactivating peptide with anti-fungal activity from sponge gourd (*Luffa cylindrica*) seeds. *Peptides*, 23(6):1019-24
- Muthumani P, Meera R, Subin M, Devi P, Kameswari B, Priya B. (2010). Phytochemical screening and anti-inflammatory, bronchodilator and antimicrobial activities of the seeds of *Luffa cylindrica*. *Res J Pharmaceut, Biol Chemical Science*, 1 (4):11-22
- Ng Y-M, Yang Y, Sze K-H, Zhang X, Zheng Y-T, Shaw P-C. (2011). Structural characterization and anti-HIV-1 activities of arginine/glutamate-rich polypeptide Luffin P1 from the seeds of sponge gourd (*Luffa cylindrica*). *J Struct Biology*, 174(1):164-72
- Kao T, Huang C, Chen B. (2012). Functional components in *Luffa cylindrica* and their effects on anti-inflammation of macrophage cells. *Food Chemistry*, 135 (2):386-95.
- Farag MA, Rasheed DM, Kamal IM. (2015). Volatiles and primary metabolites profiling in two *Hibiscus*



- sabdariffa (roselle) cultivars via headspace SPME-GC-MS and chemometrics. *Food Res International*, 78:327–35
- Farag MA, Wessjohann LA. (2012). Volatiles profiling in medicinal licorice roots using steam distillation and solid-phase microextraction (SPME) coupled to chemometrics. *Journal of Food Science*, 77(11):C1179–84
- Smith CA, Want EJ, O'Maille G, Abagyan R, Siuzdak G. (2006). GCMS: processing mass spectrometry data for metabolite profiling using nonlinear peak alignment, matching, and identification. *Analytical Chemistry*, 78(3):779–87.
- Fan W, Qian MC. (2006). Identification of aroma compounds in Chinese 'Yanghe Daqu' liquor by normal phase chromatography fractionation followed by gas chromatography [sol] olfactometry. *Flavour Fragrance Journal*, 21(2):333–42.
- Park S-N, Lim YK, Freire MO, Cho E, Jin D, Kook J-K. (2012). Antimicrobial effect of linalool and a-terpineol against periodontopathic and cariogenic bacteria. *Anaerobe*, 18(3):369–72.
- Zhang Z-Z, Li Y-B, Qi L, Wan X-C. (2006). Antifungal activities of major tea leaf volatile constituents toward *Colletorichum camelliae* Masea. *Journal of Agriculture Food Chemistry*, 54(11):3936–40
- Griego FY, Bogen KT, Price PS, Weed DL. (2008). Exposure, epidemiology and human cancer incidence of naphthalene. *Regul Toxicol Pharmaceuticals*, 51 (2):22–6
- Shmuel Y. Dictionary of food compounds with CD-ROM, 2nd ed: CRC Press, Taylor and Francis Group; 2012. 2346 p
- Zhou J, Xie G, Yan X. Encyclopedia of Traditional Chinese Medicines - Molecular Structures, Pharmacological Activities, Natural Sources and Applications: Springer-Verlag Berlin Heidelberg; 2011. 730 p
- Tang W, G. E.(1992) *Luffa cylindrica* (L.) Roem. In: Chinese Drugs of Plant Origin: Springer, Berlin, Heidelberg.
- Tanaka S, Uno C, Akimoto M, Tabata M, Honda C, Kamisako W. (1991). Anti-allergic effect of bryonolic acid from *Luffa cylindrica* cell suspension cultures. *Planta Med*, 57(6):527–30
- Urbanová T, Tarkowská D, Novák O, Hedden P, Strnad M. (2013). Analysis of gibberellins as free acids by ultra performance liquid chromatography– tandem mass spectrometry. *Talanta* 2013;112:85–94.
- Amin A, Gali-Muhtasib H, Ocker M, Schneider-Stock R. (2009). Overview of major classes of plant-derived anticancer drugs. *Int J Biomedical Sciences*, 5(1):1-11
- Ediriweera MK, Tennekoon KH, Samarakoon SR. (2019). In vitro assays and techniques utilized in anticancer drug discovery. *Journal of Applied Toxicology*, 39(1):38- 71.
- Al-Fatlawi AAY, Al-Salih ARH, Yassen MAR. (2017). β sitosterol protects against cisplatin-induced nephrotoxicity through amelioration of oxidative stress in rats. *Muthanna Medical Journal*, 4(2):60-74
- Cárdeno A, Aparicio-Soto M, Montserrat-de la Paz S, Bermudez B, Muriana FJ, Alarcón-de-la-Lastra C. (2015). Squalene targets pro-and anti-inflammatory mediators and pathways to modulate over-activation of neutrophils, monocytes and macrophages. *Journal of Functional Foods*, 14:779-790.
- Patel K, Gadewar M, Tahilyani V, Patel D.K. (2013). A review on pharmacological and analytical aspects of diosmetin: a concise report. *Chinese journal of integrative medicine*, 19(10):792-800.
- Parama D, Boruah M, Kumari Y, Rana, V, Banik K, Harsha C. (2020). Diosgenin, a steroidal saponin, and its analogues: Effective therapies against different chronic diseases. *Life sciences* 2020, 118182
- Elekofehinti O.O.(2015). Saponins: Anti- diabetic principles from medicinal plants-A review. *Pathophysiology*, 22:95- 103. doi:10.1016/j.pathophys.2015.02.001
- Garai S, Ghosh R, Bandopadhyay PP, Mandal NC Chattopadhyay A. (2018). Anti-microbial and anti-cancer properties of echinocystic acid extracted from *Luffa cylindrica*. *J Food Process Technol* 2018;9:2, doi: 10.4172/57-7110.1000717
- Yadav R, Yadav BS, Yadav R.B. (2017). Phenolic profile and antioxidant activity of thermally processed sponge gourd (*Luffa cylindrica*) as studied by using high performance thin layer chromatography (HPTLC). *International Journal of Food Properties*, 20(9):2096-2112.
- Akther F, Rahman A, Proma JJ, Kabir Z, Paul PK, Rahmatullah M. (2014). Methanolic extract of *Luffa cylindrica* fruits show antihyperglycemic potential In Swiss albino mice. *Advances in Natural and Applied Sciences*, 8(8):62-65.
- Surana AR, Kumbhare MR, Wagh R.D. (2016). Estimation of total phenolic and total flavonoid content and

- assessment of in vitro antioxidant activity of extracts of *Hamelia patens* Jacq. stems. *Research Journal of Phytochemistry*. 10(2):67- 74.
- He J, Wu ZY, Zhang S, Zhou Y, Zhao F, Peng ZQ, Hu ZW. (2014). Optimisation of microwave-assisted extraction of tea saponin and its application on cleaning of historic silks. *Journal of Surfactants and Detergents*, 17(5):919-28.
- Adeniyi SA, Orjiekwe CL, Ehiagbonare J.E.(2009). Determination of alkaloids and oxalates in some selected food samples in Nigeria. *African Journal of Biotechnology*, 8(1):110-2.
- Alagbe, J.O., Agubosi, O.C.P., Oluwafemi, R.A and Gabriel Zakara (2022). Efficacy of *Trichilia monadelpha* stem bark on the growth performance of growing rabbits. *British Journal of Innovation in Science, Research and Development*, 1(2): 10-19.
- Alagbe, J.O. (2022). *Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: haematology and serum biochemical indices. *Central Asian Journal of Theoretical and Applied Sciences* 3(2): 19-29.
- Alagbe, J.O., Adediji, M.O., Habiba, Z., Nwosu, Gloria and Wyedia Dabara Comfort (2021). Physico-chemical properties of *Indigofera zollingeriana* seed oil. *Asian Journal of Advances in Medical Science* 3(4): 306-308.
- Alagbe, J.O., Shittu, M.D and Ushie, F.T. (2021). GC-MS analysis of methanolic stem bark extract of *Zollingeriana indigofera*. *Asian Journal of Advances in Research* 11(4): 144-146.
- Alagbe, J.O (2021). Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phyto-genic Feed Additive in Growing Rabbit Diets: Growth Performance and Caecal Microbial Population. *Concept in Dairy and Veterinary Sciences*. 4(2):2021.
- Adeyale, A.O., Alagbe, J.O., Adeoye, Adekemi. O. (2021). Dietary Supplementation of *Rauvolfia Vomitoria* Root Extract as A Phyto-genic Feed Additive in Growing Rabbit Diets: Haematology and serum biochemical indices. *International Journal of Orange Technologies*, 3(3): 1-12.
- Singh, A.S., Alagbe, J.O., Sharma, S., Oluwafemi, R.A and Agubosi, O.C.P. (2021). Effect of dietary supplementation of melon (*Citrullus linatus*) seed oil on the growth performance and antioxidant status of growing rabbits. *Journal of Multidimensional Research and Reviews*, 2(1): 78-95.
- Shittu, M.D., Alagbe, J.O., Adejumo, D.O., Ademola, S.G., Abiola, A.O., Samson, B.O and Ushie, F.T. (2021). Productive Performance, Caeca Microbial Population and Immune-Modulatory Activity of Broiler Chicks Fed Different Levels *Sida Acuta* Leaf Extract in Replacement of Antibiotics. *Bioinformatics and Proteomics Open Access Journal* 5(1): 000143.
- Alagbe, J.O. (2021). *Prosopis africana* stem bark as an alternative to antibiotic feed additives in broiler chicks diets: Performance and Carcass characteristics. *Journal of Multidimensional Research and Reviews*, 2(1): 64-77.
- Alagbe, J.O. (2021). *Daniellia oliveri* leaf extracts as an alternative to antibiotic feed additives in broiler chicken diets: Meat Quality and Fatty acid composition. *Indonesian Journal of Innovation and Applied Sciences* 1(3): 177-186.
- Alagbe, J.O., Adeoye, Adekemi and Oluwatobi, O.A. (2020). Proximate and mineral analysis of *Delonix regia* leaves and roots. *International Journal on Integrated Education*. 3(10): 144-149.
- Musa, B., Alagbe, J.O., Adegbite Motunrade Betty, Omokore, E.A. (2020). Growth performance, caeca microbial population and immune response of broiler chicks fed aqueous extract of *Balanites aegyptiaca* and *Alchornea cordifolia* stem bark mixture. *United Journal for Research and Technology*, 2(2):13-21.
- Shittu, M.D and Alagbe, J.O. (2020). Phyto-nutritional profiles of broom weed (*Sida acuta*) leaf extract. *International Journal of Integrated Education*. 3(11): 119-124
- Alagbe, J.O., Shittu, M.D and Eunice Abidemi Ojo (2020). Prospect of leaf extracts on the performance and blood profile of monogastric – A review. *International Journal of Integrated Education*. 3(7): 122-127.
- Alagbe, J.O (2019). Proximate, mineral and phytochemical analysis of *Piliostigma thonningii* stems bark and roots. *International Journal of Biological, Physical and Chemical Studies*, 1(1): 1-7.
- Alagbe, J.O. (2019). Haematology, serum biochemistry, relative organ weight and bacteria count of broiler chicken given different levels of *Luffa aegyptiaca* leaf extracts. *International Journal of Advanced Biological and Biomedical Research*. 7(4):382-392.
- Alagbe, J.O and Omokore, E.A. (2019). Effect of replacing soya bean meal with *Indigofera zollingeriana* leaf meal on the performance and carcass characteristics of growing rabbits. *International Journal of Multidisciplinary Research and Development*. 6(5): 74-77.