



Bioactive compounds of *Prosopis Africana* oil (African mesquite) using gas chromatography and mass spectrometry (GC-MS) technique

Alagbe John Olujimi^{a*}

a. Department of Animal Science, University of Abuja, Nigeria

Abstract: Phytochemicals (bioactive compounds) are generally regarded as chemicals of plant origin used by plants for growth, defense against pathogens, and other competitors. They also have a wide range of pharmacological activities, including antiarrhythmic, antioxidant, antimalarial, vasodilatory, antimicrobial, analgesics, hepato-protective, anti-inflammatory, hypo-cholesterolemic, anti-androgenic, antiviral, antifungal, and immune-modulatory. This study was carried out to investigate the bioactive compounds of *Prosopis africana* oil (African mesquite) using the gas chromatography–mass spectrometry technique. A total of 73 bioactive compounds (77.16 %) were identified based on their peak areas. The major compounds identified in the oil sample were prospering A (29.90 %), caryophyllene (12.33 %), 2, 4-bis (1-phenylmethyl) phenol (5.80 %), gallic acid (5.22 %), β -cycloidal (3.11 %), β -sitosterol (2.50 %), α -terpinene (2.09 %), and ellagic acid (2.04 %). In conclusion, *Prosopis africana* oil is loaded with phytochemicals and has several health-promoting properties, and it can also be used as a remedy for the treatment of several ailments in human beings and animals.

Keywords: Phytochemicals, *Prosopis Africana*, Pharmacological properties, Gas chromatography

1. Introduction

Prosopis africana (African mesquite) is an evergreen multipurpose leguminous plant belonging to the family Fabaceae. It has about 45 species, and it is widely distributed in most of Africa, Asia, and America ([Adikwu et al., 2001](#); [Ugwoke et al., 2019](#)). The seeds of *Prosopis africana* are rich in phyto-constituents such as alkaloids, tannins, flavonoids, terpenoids, steroids, saponins, glycosides, and phenols with significant therapeutic properties ([Alagbe et al., 2021](#); [Olorunmaiye et al., 2019](#)). The seeds have antioxidant, antifungal, antimicrobial, antifungal, angelic, antipyretic, anti-proliferative, anti-inflammatory, hepato-protective, antiprotozoal, anti-androgenic, and cytotoxic properties ([Harzallah-Skhirri & BenJannet, 2005](#); [Sharma et al., 2022](#)), and they also contain important minerals such as calcium, potassium, zinc, manganese, magnesium, copper, and sodium ([Ferguson et al., 2005](#); [Oluwafemi et al., 2021](#)). These minerals are vital for the activation of key enzymes that provide greater protection from diseases ([Alagbe, 2022b](#)).

Moreover, *Prosopis africana* seeds contain amino acids alanine, proline, valine, and isoleucine, which are the main structural proteins found in the connective tissues of animals ([Barminas et al., 1998](#); [Sirmah et al., 2009](#)). The medicinal properties of plants depend on their composition of phyto-constituents or bioactive compounds ([Ogunshe et al., 2007](#)). The stem bark and root decoction of *Prosopis* are used traditionally for the treatment of rheumatism, cough, malaria, bronchitis, skin diseases, and gastrointestinal and sexually transmitted infections. Fermented seeds of the plant can be used in the preparation of soup because of their sweet smell ([Gberikon et al., 2015](#); [Platt, 1964](#)).

Preclinical investigations have shown the pharmacological responses of *Prosopis africana* oil to inhibit the activities of some pathogenic organisms, such as *Salmonella* spp, *Candida albicans*, *Escherichia coli*, *Bacillus subtilis*, *Bacillus pumilus*, *Bacillus licheniformis*, and *Bacillus megaterium* ([Alagbe, 2022a](#); [Oguntoyinbo et al., 2010](#)). Aqueous and ethanolic extracts from the leaves and stem bark of *Prosopis africana* can be used for the treatment of diarrhea and coccidiosis in poultry ([Adewale et al., 2021](#)). Identifying the bioactive compounds in

[Received] 14 April 2023; Accepted 23 June 2023; Published (online) 27 June 2023]

Finesse Publishing stays neutral regarding jurisdictional claims published maps.



Attribution 4.0 International (CC BY 4.0)

Corresponding email: demsonfarms@yahoo.com (Alagbe, J.O)

DOI: 10.61363/fsamr.v2i1.60

a medicinal plant could lead to the production of novel drugs (Alagbe, 2020). One key method in identifying unknown compounds, quantification, and characterization is the use of gas chromatography–mass spectrometry (GC-MS) technique.

Investigating the bioactive compounds in a sample will give a clue about its efficacy and safety. Hence, given the immense medicinal properties of the *Prosopis africana* seed. This study was designed to evaluate the phytochemicals or bioactive compounds in *Prosopis africana* oil using the GC-MS technique.

2. Materials and methods

2.1. Experimental site

This study was carried out at the Department of Animal Science Laboratory, University of Abuja main campus along Airport Road, Gwagwalada, Abuja, Nigeria, located between latitudes 8o571 and 8o551N and longitude 7o051 and 7o061E (Balogun & Balogun, 2001).

2.2. Collection, identification, and processing of *Prosopis africana* oil

Prosopis africana seeds were purchased from a local market in Gwagwalada, Nigeria. It was identified and authenticated at the Herbarium of the Department of Biological Sciences, University of Abuja, Nigeria, with a voucher number ABJ/09A/20. It was shade-dried for 13 days, ground into powder using a laboratory blender (Panasonic, Model EDH-023), and stored in a labeled polythene bag for further testing.

2.3. Extraction of *Prosopis africana* oil using the steam distillation method

Prosopis africana oil was extracted using the steam-distillation technique; the entire extraction procedure requires a digital weighing scale, a round-bottom flask, distilled water, a glass yarn heating mantle, a measuring cylinder, and a separatory funnel. 50 grams of ground *Prosopis* seed were weighed into a round-bottom flask and mixed with 250 mL of distilled water. The mixture was placed in a glass jar with a heating mantle and adjusted to a temperature of 80 °C before setting up the condenser above the round-bottom flask. The mixture is boiled vigorously for 15 minutes, and the distillate is collected in a beaker until no more droplets of oil come over, followed by the addition of 5 grams of sodium chloride, and stirred vigorously. Thereafter, the distillate was transferred into the separatory funnel and evaporated in a steam bath to obtain *Prosopis africana* oil.

2.4. Bioactive compounds of PRAO using the GC-MS technique

Bioactive compounds of PRAO were identified using LABTRON GC-MS (GC-MS-879). It is a high-precision gas chromatography-mass spectrometer with a pre-filter mass analyzer and an electron multiplier ensuring high sensitivity. Gas chromatography (Model – GC-87) has the following technical specifications; flow rate (1-15 ml/min), maximum flow rate (10 ml/min), inlet temperature (450 °C), pressure range (0 – 100 psi), heating rate (up to 120 °C / min), room temperature (40 °C – 450 °C), pressure control mode (electronic pressure control), split mode (split / split less), split ratio (1000: 1) and temperature programming (7 stages / 8 platforms) while the mass spectrometry (Model- MS-89): mass range (1.5 – 1000 amu), ion – source temperature (100 °C – 350 °C), GC-MS interface temperature (450° C), scan rate (up to 1000 amu/sec), stability (\pm 0.10 amu/ 48 hours), filament emission current (0 - 350° C μ A), EI source ionization energy (5 eV – 250 eV), sensitivity (S/N is \geq 30:1) and vacuum (Turbo molecular pump: 67 L/s). Volatile compounds were identified with standard compounds in the National Institute of Standards and Technology mass spectral library using the Chemstation library database. The percentage area was based on the ratio between the peak area of each compound and the sum of the peak areas of all compounds.

Table 1: Bioactive compounds of *Prosopis Africana* oil as detected by GC-MS

Peak No	RT (min)	Area (%)	Name of compound
1	2.77	29.90	Prosogerin A (6 Methoxy-7-hydroxyl dioxyflavone)
2	3.93	5.80	2,4-bis (1-phenylmethyl) phenol
3	4.03	0.63	β -phenethylamine
4	4.51	0.14	2,4,6-tris(1-phenylmethyl) phenol
5	4.92	0.16	α -pinene
7	5.22	2.09	α -terpinene
8	5.91	0.46	β -pinene
9	6.07	0.12	β -myrcene



10	6.34	0.74	α -phellandrene
11	6.88	0.11	α -terpinolene
12	7.22	0.35	γ -terpinene
13	8.09	0.03	1-terpineol
14	8.44	0.07	4-terpineol
15	9.57	0.18	Humulene
16	9.88	12.33	Caryophyllene
17	10.12	1.85	Copaene
18	11.55	0.21	Cis-linalolide
19	11.83	0.30	α -Selinene
20	11.90	0.06	γ -Elemene
21	12.10	0.01	α -Gurjunene
22	12.33	0.42	β -Elemene
23	12.68	3.11	β -Cyclocitral
24	12.75	0.05	3-hexenyl-2-methylbutanoate
25	12.92	0.02	Exo-methyl-camphenilol
26	13.10	0.06	Caryophyllene oxide
27	13.47	0.03	Benzene -1-methoxy-2-methyl
28	13.55	0.18	Napthalene, 1,2 hydro-1,1, 6 -trimethyl
29	13.80	1.55	Cycloheptasiloxane, tetradecamethyl
30	13.87	1.41	Nerolidyl acetate
31	13.91	0.001	1-Cyclohexene-1- butanal, alpha, 2, 6, 6- tetramethyl
32	14.00	0.11	Cyclooctasiloxane, hexadecamethyl
33	14.31	0.60	2,2, 6-Trimethyl-2H, 5Hpyrano(3,2-c) quinoline-5one
34	14.49	0.27	Dibutyl phthalate
35	14.52	0.70	Acetic acid, dodecyl ester
36	14.70	0.64	1,1, 5-Trimethyl-1,2-dihydro naphthalene
37	14.83	0.02	Alloaromedendrene oxide
38	14.92	0.03	Butylated Hydroxytoluene
39	14.95	0.01	Octacosane
40	16.71	0.16	Kaur-16-ene
41	16.37	0.10	1,5,8-p-Menthatriene
42	16.55	0.20	Spicigerine
43	18.28	0.07	Prosophylline
44	18.84	2.04	Ellagic acid
45	18.92	0.55	Apigenin-6-glucoside
46	19.20	1.80	Quercetin-3-rhamnoside
47	19.44	0.01	Isoprosopilosine
48	19.75	1.50	Prosopilosidine
49	19.87	0.02	Trimethyl-tetrahydronaphtalene
50	21.21	0.21	Spathulenol
51	21.40	0.01	1,3-benzodioxole,5-(2,2-dimethyl)
52	21.61	0.22	2-nonen-4-one
53	22.02	0.11	Bicyclo [2.2.1] heptan-3-one
54	25.04	2.50	β -sitosterol
55	25.80	0.32	Cis-vaccenic acid
56	26.11	0.16	3-chloro-N-isochron-1-methyl propionamide
57	26.70	0.03	3,4 - dichloro benzonitrile
58	26.84	0.07	Mannitol, 1,4-di-O-methyl tetraacetate
59	27.02	0.14	β -1-rhamnofuranoside thio-octyl
60	27.18	0.03	Guaia-1(10), 11-diene
61	27.35	0.12	3,4 - dimethyl phenyl heptyl ether
62	27.80	0.09	1,1,5 -trimethyl -1,2 -dihydro naphthalene
63	27.93	0.03	1-cyclohexane-1-propanal

64	28.06	0.04	1-Trimethylsilylpent-1-en-4-one
65	28.93	0.02	β -ocimene
66	28.98	0.05	Trans- β -ocimene
67	33.40	0.18	1-Methyl-6-methylene bicyclo heptane
68	33.73	0.04	1,6,10,14 -Hexadecatetraen-3-ol
69	33.81	0.01	Alloaromadendrene oxide (1)
70	33.92	0.40	1-cyclohexane-1-propanal
71	36.27	0.18	1-Formyl -2,2,6 -trimethyl cyclohexane
72	36.84	5.22	Gallic acid
73	36.93	0.06	6-Isopropenyl-4,8-dimethyl naphthalene-2-ol
74	37.10	0.01	1,6,10-Dodecatrien-3-trimethyl -ol
Total		77.16	

RT: Retention time (min)

3. Results and discussion

3.1. Bioactive compounds of *Prosopis africana* oil

The bioactive compounds of *Prosopis africana* oil are presented in Table 1. A total of 73 bioactive compounds (77.16 %) were identified based on their peak areas. The major compounds identified in the oil sample were prospering A (29.90 %), caryophyllene (12.33 %), 2, 4-bis (1-phenylmethyl) phenol (5.80 %), gallic acid (5.22 %), β -cycloidal (3.11 %), β -sitosterol (2.50 %), α -terpinene (2.09 %), and ellagic acid (2.04 %). Other minor compounds includes; β -phenethylamine (0.63 %), 2, 4, 6-tris (1-phenylethyl) phenol (0.14 %), α -pinene (0.16 %), β -myrcene (0.12 %), α -phellandrene (0.74 %), α -terpinolene (0.11 %), γ -terpinene (0.35 %), 1-terpineol (0.03 %), 4-terpineol (0.07 %), humulene (0.18 %), copaene (0.85 %), cis-linaloxide (0.21 %), α -selinene (0.30 %), γ -elemene (0.06 %), α -gurjunene (0.01 %), β -elemene (0.42 %), 3-hexenyl-2-methylbutanoate (0.05 %), exo-methyl-camphenilol (0.02 %), benzene -1-methoxy-2-methyl (0.03 %), naphthalene, 1,2 hydro-1,1, 6 -trimethyl (0.18 %), cycloheptasiloxane, tetradecamethyl (1.55 %), nerolidyl acetate (1.41 %), 1-cyclohexene-1- butanal, alpha, 2, 6, 6- tetramethyl (0.001 %), cyclooctasiloxane, hexadecamethyl (0.11 %), 2,2, 6-trimethyl-2H, 5H pyrano (3,2-c) quinolin-5-one (0.60 %), dibutyl phthalate (0.27 %), acetic acid, dodecyl ester (0.70 %), 1,1, 5-trimethyl-1,2-dihydronaphthalene (0.64 %), alloaromedendrene oxide (0.02 %), butylated hydroxytoluene (0.03 %), octacosane (0.01 %), kaur-16-ene (0.16 %), 1,5,8-p-menthatriene (0.10 %), spicigerine (0.20 %), prosophylline (0.07 %), apigenin-6-glucoside (0.55 %), quercetin-3-rhamnoside (1.80 %), isoprosopilosine (0.01 %), prosopilosidine (1.50 %), trimethyl-tetrahydronaphtalene (0.02 %), spathulenol (0.21 %), 1,3-benzodioxole,5-(2,2-dimethyl) (0.01 %), 2-nonen-4-one (0.22 %), bicyclo [2.2.1] heptan-3-one (0.11 %), cis-vaccenic acid (0.32 %), 3-chloro-N-isochroman-1-methyl propionamide (0.16 %), 3,4 - dichlorobenzonitrile (0.03 %), mannitol, 1,4-methyl tetraacetate (0.07 %), β -1-rhamnofuran oside thio-octyl (0.14 %), guaia-1(10), 11-diene (0.03 %), 3,4 - dimethyl phenyl heptyl ether (0.12 %), 1,1,5 -trimethyl -1,2 -dihydronaphthalene (0.09 %), 1-cyclohexane-1-propanal (0.03 %), 1-trimethylsilylpent-1-en-4-yne (0.04 %), β -ocimene (0.02 %), trans- β -ocimene (0.05), -methyl-6-methylenebicycloheptane (0.18 %), 1,6,10,14 -hexadecatetraen-3-ol (0.04 %), alloaromadendrene oxide (1) (0.01 %), 1-cyclohexane-1-propanal (0.40 %), 1-formyl -2,2,6 -trimethyl cyclohexane (0.18 %), 6-isopropenyl-4,8-dimethyl nephthalen-2-ol (0.06 %) and 1,6,10-dodecatrien-3-trimethyl -ol (0.01 %). However, all these compounds have significant pharmacological effects ([Sharma et al., 2022](#); [Shittu et al., 2022](#)).

Prosogerin A has been reported to possess antimicrobial and anti-inflammatory properties ([Agubosi et al., 2021](#)). Caryophyllene occurs naturally in essential oils of several plants, including *Piper nigrum*, *Lantana camara* leaf oil, and *Capaifera* spp, and has been reported to function as an antioxidant, thus scavenging free radicals ([Pagare et al., 2015](#)). 2,4-bis (1-phenylethyl) phenol, α -terpinene, 2,4,6-tris(1-phenylethyl) phenol, β -myrcene, α -terpinolene, and α -phellandrene had been reported by researchers to have anti-inflammatory and hypocholesterolemic properties ([Olasan, 2016](#); [Silva et al., 2015](#)). Humulene, β -sitosterol, α -gurjunene, α -selinene, and exo-methyl-camphenilol possess hepato-protective, anti-proliferative, antifungal, antipyretic, and antihelminthic properties ([Alagbe, 2023](#); [Oluwafemi et al., 2021](#); [Shibula & Velavan, 2015](#)). Naphthalene, 1,2 hydro-1,1, 6 - trimethyl, and benzene -1-methoxy-2-methyl can perform nematocidal and anti-androgenic activities ([Borokini & Omotayo, 2012](#); [Dua et al., 2010](#)). The other bioactive compounds in lower concentrations contained in *Prosopis Africana* oil also have significant therapeutic properties and are used in ayurvedic remedies for a variety of ailments ([Ayoola et al., 2008](#)). However, the result obtained in this experiment agrees with the findings of [Lokesh et al. \(2018\)](#); [Nakano et al. \(2002\)](#), and [Aneela et al. \(2014\)](#) but contrary to the reports



of [Agubosi et al. \(2021\)](#). These differences are influenced by factors like species, climate, harvest stage, location, storage condition, and method of extraction of the oil ([Idris et al., 2009](#)).

4. Conclusion

It can be concluded that *Prosopis Africana* oil possesses several health-promoting properties (antimicrobial, antifungal, antiviral, anti-fibrotic, immune-modulatory, cytotoxic, antipyretic, antitumor, antihelminthic, and hepato-protective) due to the presence of bioactive compounds. These compounds can reduce the rising concern about environmental pollution, antibiotic resistance, and food safety. It could also enhance livestock performance and the production of novel drugs.

CRedit authorship contribution statement

Alagbe John Olujimi, Experiment, analysis of data, writing-original draft, editing, and proofreading.

Funding

This work was not financially supported by any funding agency.

Declaration of competing interest

The authors declare no conflict of interest.

Acknowledgments

The author(s) acknowledge that they did not receive any financial support from any funding agency, whether public, private, or non-profit. The study was conducted independently without external funding.

References

- Adewale, A., Alagbe, J., & Adeoye, A. O. (2021). Dietary supplementation of *Rauvolfia vomitoria* root extract as a phyto-genic feed additive in growing rabbit diets: Haematology and serum biochemical indices. *International Journal of Orange Technologies*, 3(3), 1-12.
- Adikwu, M., Ezeabasili, S., & Esimone, C. (2001). Evaluation of the physico-chemical properties of a new polysaccharide gum from *Prosopis africana*. *Bollettino Chimico Farmaceutico*, 140(1), 40-45.
- Agubosi, O., Oluwafemi, R., & Alagbe, J. (2021). The effect of processing on the proximate, mineral and vitamin composition of Neem leaves (*Azadirachta indica*) grown in Gwagwalada, FCT, Abuja. *Abuja Journal of Agriculture and Environment*, 1(1), 293-299.
- Alagbe, J. (2020). Chemical evaluation of proximate, vitamin and amino acid profile of leaf, stem bark and root of *Indigofera Tinctoria*. *European Journal of Research Development and Sustainability*, 1(1), 5-12.
- Alagbe, J. (2022a). Characterization Of Bioactive Compounds in *Luffa Aegyptiaca* Leaf Ethanolic Extracts Using Gas Chromatography And Mass Spectrometer: Characterization Of Bioactive Compounds in *Luffa Aegyptiaca* Leaf Ethanolic Extracts Using Gas Chromatography And Mass Spectrometer. *Food Science & Applied Microbiology Reports*, 1(2), 21-28.
- Alagbe, J. (2022b). *Prosopis africana* (African mesquite) oil as an alternative to antibiotic feed additives on broiler chickens diets: haematology and serum biochemical indices. *Central Asian Journal of Theoretical and Applied Science*, 3(2), 19-29.
- Alagbe, J., Adedeji, M., Habiba, Z., Nwosu, G., & Comfort, W. D. (2021). Physico-chemical properties of *Indigofera zollingeriana* seed oil. *Asian Journal of Advances in Medical Science*, 3(4), 306-308.
- Alagbe, J. O. (2023). Bioactive compounds in ethanolic extract of *Strychnos innocua* root using gas chromatography and mass spectrometry (GC-MS). *Drug Discovery*, 17, e4dd1005.
- Aneela, S., Dey, A., & De, S. (2014). GC-MS analysis of methanolic extract of *Prosopis spicigera*. *International Journal of Phytopharmacology*, 5(3), 168-171.
- Ayoola, G., Coker, H., Adesegun, S., Adepoju-Bello, A., Obaweya, K., Ezennia, E. C., & Atangbayila, T. (2008). Phytochemical screening and antioxidant activities of some selected medicinal plants used for malaria therapy in Southwestern Nigeria. *Tropical Journal of Pharmaceutical Research*, 7(3), 1019-1024.
- Balogun, O., & Balogun, O. Y. (2001). *The Federal Capital Territory of Nigeria: a geography of its development*. Ibadan university press.
- Barminas, J., Maina, H., & Ali, J. (1998). Nutrient content of *Prosopis africana* seeds. *Plant Foods for Human Nutrition*, 52(4), 325-328.

- Borokini, T. I., & Omotayo, F. O. (2012). Phytochemical and ethnobotanical study of some selected medicinal plants from Nigeria. *Journal of Medicinal Plants Research*, 6(7), 1106-1118.
- Dua, V., Pandey, A., & Dash, A. (2010). Adulticidal activity of essential oil of Lantana camara leaves against mosquitoes. *Indian Journal of Medical Research*, 131(3), 434-439.
- Ferguson, L. R., Zhu, S. t., & Harris, P. J. (2005). Antioxidant and antigenotoxic effects of plant cell wall hydroxycinnamic acids in cultured HT-29 cells. *Molecular nutrition & food research*, 49(6), 585-593.
- Gberikon, G., Agbulu, C., & Yaji, M. (2015). Nutritional composition of fermented powdered Prosopis africana soup condiment with and without inocula.
- Harzallah-Skhiri, F., & BenJannet, H. (2005). Flavonoids diversification in organs of two prosopis farcta (banks & sol.) eig.(leguminosea, mimosoideae) populations occurring in the northeast and the southeast of Tunisia. *J Appl Sci Res*, 1(2), 130-136.
- Idris, S., Ndukwe, G., & Gimba, C. (2009). Preliminary phytochemical screening and antimicrobial activity of seed extracts of Persea americana (avocado pear).
- Lokesh, K., Sonal, D., Pradeep, P., & Dobhai, M. (2018). Phytochemicals from Prosopis species. Suresh Gyan Vihar University International Journal of Environment. *Science and Technology*, 4(1), 9-32.
- Nakano, H., Fujii, Y., Yamada, K., Kosemura, S., Yamamura, S., Hasegawa, K., & Suzuki, T. (2002). Isolation and identification of plant growth inhibitors as candidate (s) for allelopathic substance (s), from aqueous leachate from mesquite (Prosopis juliflora (Sw.) DC.) leaves. *Plant Growth Regulation*, 37(2), 113-117.
- Ogunshe, A. A., Omotosho, M. O., & Ayansina, A. (2007). Microbial studies and biochemical characteristics of controlled fermented afiyo-a Nigerian fermented food condiment from Prosopis africana (Guill and Perr.) Taub. *Pakistan Journal of Nutrition*, 6(6), 620-627.
- Oguntoyinbo, F. A., Huch, M., Cho, G.-S., Schillinger, U., Holzapfel, W. H., Sanni, A. I., & Franz, C. M. (2010). Diversity of Bacillus species isolated from okpehe, a traditional fermented soup condiment from Nigeria. *Journal of Food Protection*, 73(5), 870-878.
- Olasan, O. (2016). Phytochemical screening of Xylopia aethiopica with emphasis on its medicinally active principles. *Journal of Medicinal Plants Research*.
- Olorunmaiye, K., Apeh, L., Madandola, H., & Oguntoye, M. (2019). Proximate and Phytochemical Composition of African Mahogany (Afzelia africana) seed and African mesquite (Prosopis africana) pod. *Journal of Applied Sciences & Environmental Management*, 23(2).
- Oluwafemi, R., Agubosi, O., & Alagbe, J. (2021). Proximate, minerals, vitamins and amino acid composition of Prosopis africana (African mesquite) seed oil. *Asian Journal of Advances in Research*, 11(1), 21-27.
- Pagare, S., Bhatia, M., Tripathi, N., Pagare, S., & Bansal, Y. (2015). Secondary metabolites of plants and their role: Overview. *Current trends in biotechnology and pharmacy*, 9(3), 293-304.
- Platt, B. (1964). Biological ennoblement: improvement of the nutritive value of foods and dietary regimens by biological agencies.
- Sharma, S., John, A. O., Xing, L., Ram, S., & Amita, K. (2022). Comparative analysis of ethanolic Juniperus thurifera leaf, stem bark and root extract using gas chromatography and mass spectroemetry. *International Journal of Agriculture and Animal Production*, 2(6), 18-27.
- Shibula, K., & Velavan, S. (2015). Determination of phytocomponents in methanolic extract of Annona muricata leaf using GC-MS technique. *International Journal of Pharmacognosy and Phytochemical Research*, 7(6), 1251-1255.
- Shittu, M., Alagbe, J. O., Ojebiyi, O. O., Ojediran, T. K., & Rafiu, T. A. (2022). Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of Polyalthia longifolia leaf extract in place of conventional antibiotics. *Animal Science and Genetics*, 18(2).
- Silva, T., Suffredini, I., Ricci, E., Fernandes, S., Gonçalves Jr, V., Romoff, P., Lago, J., & Bernardi, M. (2015). Antinociceptive and anti-inflammatory effects of Lantana camara L. extract in mice. *Revista Brasileira de plantas medicinais*, 17, 224-229.
- Sirmah, P., Dumarçay, S., Masson, E., & Gérardin, P. (2009). Unusual amount of (-)-mesquitol from the heartwood of Prosopis juliflora. *Natural Product Research*, 23(2), 183-189.
- Ugwoke, A., Oluka, S., & Eze, P. (2019). PHYSICOMECHANICAL PROPERTIES OF IRON TREE SEED (PROSOPIS AFRICANA) FROM SOUTH EASTERN NIGERIA. *Journal of Agricultural Engineering and Technology (JAET)*, 24(1), 43-50.